PB89-201875

Pentachlorophenol: Health and Environmental Effects Profile

(U.S.) Environmental Protection Agency, Cincinnati, OH

1986

U.S. Department of Commerce
National Technical Information Service
NTIS

TECHNICAL REPORT DATA
(Please read Instructions on the reverse before complet

1. REPORT NO. EPA/600/J-86/503	2.	3. PB89-201875
4. TITLE AND SUBTITLE Pentachlorophenol: Health and Environmental Effects Profile		5. REPORT DATE 6. PERFORMING ORGANIZATION CODE
7. AUTHOR(S) H. Choudhury, J. Coleman, C.T. DeRosa and J.F. Stara		8. PERFORMING ORGANIZATION REPORT NO.
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT NO. 11. CONTRACT/GRANT NO.
12. SPONSORING AGENCY NAME AND ADDRESS Environmental Criteria and Assessment Office Office of Research and Development U.S. Environmental Protection Agency Cincinnati, OH 45268		13. TYPE OF REPORT AND PERIOD COVERED 14. SPONSORING AGENCY CODE EPA/600/22

15. SUPPLEMENTARY NOTES

Toxicology and Industrial Health 2(4):483-571 (1986)

16. ABSTRACT

Pentachlorophenol is used as an industrial wood preservative for utility poles, crossarms, fence posts, and other purposes(79%); for NaPCP (12%); and miscellaneous including mill uses, consumer wood preserving formulations and herbicide intermediate (9%). Photolysis and microbial degradation are the important chemical removal mechanisms for pentachlorophenol (PCP) in water. Direct photolysis may be an important environmental sink for PCP present in the atmosphere. Soil degradation studies indicate the PCP is biodegradable; microbial decomposition is an important and potentially dominant removal mechanism in soil. Monitoring studies have confirmed the widespread occurrence of PCP in surface waters, groundwater, drinking water and industrial effluents. Acute toxicity data indicated that salmonids are more sensitive to the toxic effects of PCP than other fish species. PCP is readily absorbed from the gastrointestinal tract of rats, mice, monkeys and humans. The primary route of excretion after oral administration of all species studied is in the urine. Teratogenicity/reproductive studies indicate PCP is fetotoxic in rats. Subchronic and chronic toxicity studies show adverse effects in the liver, kidney and immune system. An ADI of 0.03 mg/kg/day or 2.1 mg/day for a 70 kg human was derived from a chronic dietary study in rats. An RQ of 100 was derived based on fetotoxic effects in rats. Inadequate evidence exists for carcinogenicity.

17. KEY WORDS AND DOCUMENT ANALYSIS

a. DESCRIPTORS	b. IDENTIFIERS/OPEN ENDED TERMS	c. COSATI Field/Group

REPRODUCED BY
U.S. DEPARTMENT OF COMMERCE
NATIONAL TECHNICAL INFORMATION SERVICE
SPRINGFIELD, VA. 22161

18. DISTRIBUTION STATEMENT Public	19. SECURITY CLASS *(This Report)* Unclassified	21. NO. OF PAGES 91
	20. SECURITY CLASS *(This page)* Unclassified	22. PRICE

EPA Form 2220-1 (Rev. 4-77) PREVIOUS EDITION IS OBSOLETE

EPA/600/J-86/503

PENTACHLOROPHENOL: HEALTH AND ENVIRONMENTAL EFFECTS PROFILE

H. CHOUDHURY,* J. COLEMAN,† C.T. DE ROSA* AND J.F. STARA*

***Environmental Criteria and Assessment Office**
Office of Health and Environmental Assessment
U.S. Environmental Protection Agency
Cincinnati, Ohio

†Center for Chemical Hazard Assessment
Syracuse Research Corporation
Syracuse, New York

NTIS is authorized to reproduce and sell this report. Permission for further reproduction must be obtained from the copyright owner.

Pentachlorophenol is used as an industrial wood preservative for utility poles, crossarms, fence posts, and other purposes (79%); for NaPCP (12%); and miscellaneous, including mill uses, consumer wood preserving formulations and herbicide intermediate (9%) (CMR, 1980). As a wood preservative, pentachlorophenol acts as both a fungicide and insecticide (Freiter, 1978). The miscellaneous mill uses primarily involve the application of pentachlorophenol as a slime reducer in paper and pulp milling and may constitute ~6% of the total annual consumption of pentachlorophenol (Crosby et al., 1981). Sodium pentachlorophenate (NaPCP) is also used as an antifungal and antibacterial agent (Freiter, 1978). Pentachlorophenol also is used as a general herbicide (Martin and Worthing, 1977).

Photolysis and microbial degradation are the important chemical removal mechanisms for pentachlorophenol in water. In surface waters, pentachlorophenol photolyzes rapidly (ECETOC, 1984; Wong and Crosby. 1981; Zepp et al., 1984); however, the photolytic rate decreases as the depth in water increases (Pignatello et al., 1983). Pentachlorophenol is readily biodegradable in the presence of accli-

1. Address correspondence to: Dr. H. Choudhury, ECAO, Office of Health and Environmental Assessment, U.S. EPA, Cincinnati, OH 45268.

2. Key words: bioaccumulation, microbial degradation, pentachlorophenol, photolysis, wood preservation.

3. Abbreviations: ADI, Acceptable daily intake; BCF, Bioconcentration factor; BOD, Biological oxygen demand; CAS, Chemical Abstract Service; CS, Composite score; DOC, Dissolved organic carbon; EC_{50}, Concentration effective to 50% of recipients; K_L, Liquid-phase mass transfer coefficient; K_{oc}, Soil sorption coefficient; K_{ow}, Octanol/water partition coefficient; LC_{50}, Concentration lethal to 50% of recipients; LD_{50}, Dose lethal to 50% of recipients; MATC, Maximum acceptable toxicant concentration; MED, Minimum effective dose; MSV, Maloney sarcoma virus; NaPCP, Sodium pentachlorophenate; NOAEL, No-observed-adverse-effect level; NOEC, No-observed-effect concentration; NOEL, No-observed-effect-level; NOLC, No-observed-lethal concentration; ppb, Parts per billion; ppm, Parts per million; RQ, Reportable quantity; RV_d, Dose-rating value; RV_e, Effect-rating value; SGOT, Serum glutamic oxaloacetic transaminase; SGPT, Serum glutamic pyruvic transaminase; STEL, Short-term exposed level; TLC, Thin-layer chromatography; TLV, Threshold limit value; UV, Ultraviolet.

mated microorganisms; however, biodegradation in natural waters requires the presence of microbes that can become acclimated. A natural river water that had been receiving domestic and industrial effluents significantly biodegraded pentachlorophenol after a 15-day lag period, while an unpolluted natural river water was unable to biodegrade the compound (Banerjee et al., 1984). Even though pentachlorophenol is in ionized form in natural waters, sorption to organic particulate matter and sediments can occur (Schellenberg et al., 1984), with desorption contributing as a continuing source of pollution in a contaminated environment (Pierce and Victor, 1978). Experimentally determined BCFs have shown that pentachlorophenol can significantly accumulate in aquatic organisms (Gluth et al., 1985; Butte et al., 1985; Statham et al., 1976; Veith et al., 1979a,b; Ernst and Weber, 1978), which is consistent with its widespread detection in fish and other organisms.

Direct photolysis may be an important environmental sink for pentachlorophenol present in the atmosphere. The detection of pentachlorophenol in snow and rain water (Paasivirta et al., 1985; Bevenue et al., 1972) suggests that removal from air by dissolution is possible.

Soil degradation studies indicate that pentachlorophenol is biodegradable; microbial decomposition is an important and potentially dominant removal mechanism in soil (Baker et al., 1980; Baker and Mayfield, 1980; Edgehill and Finn, 1983; Kirsch and Etzel, 1973; Ahlborg and Thunberg, 1980). The degree to which pentachlorophenol leaches in soil is dependent on the type of soil. In soils of neutral pH, leaching may be significant, but in acidic soils, adsorption to soil generally increases (Callahan et al., 1979; Sanborn et al., 1977). The ionized form of pentachlorophenol may be susceptible to adsorption in some soils (Schellenberg et al., 1984). In laboratory soils, pentachlorophenol decomposes faster in soils of high organic content as compared with low organic content, and faster when moisture content is high and the temperature is conducive to microbial activity. Half-lives are usually ~2-4 weeks (Crosby et al., 1981).

Monitoring studies have confirmed the widespread occurrence of pentachlorophenol in surface waters, groundwater, drinking water and industrial effluents (see Table 2). The U.S. EPA's National Urban Runoff Program and National Organic Monitoring Survey reported frequent detections in storm water runoff and public water supplies (Cole et al., 1984; Mello, 1978). Primary sources by which pentachlorophenol may be emitted to environmental waters may be through its use in wood preservation and the associated effluents and its pesticidal applications. Pentachlorophenol can be emitted to the

atmosphere by evaporation from treated wood or water surfaces, by releases from cooling towers using pentachlorophenol biocides or by incineration of treated wood (Skow et al., 1980; Crosby et al., 1981). Pentachlorophenol has been detected in ambient atmospheres (Cautreels et al., 1977), in snow and rain water (Paasivirta et al,. 1985; Bevenue et al., 1972) and in emissions from hazardous waste incineration (Oberg et al., 1985). The U.S. Food and Drug Administration's Total Diet Study (conducted between 1964 and 1977) found pentachlorophenol residues in 91/4428 ready-to-eat food composites (See Tables 4 and 5). The average American dietary intake of pentachlorophenol during 1965-1969 was estimated to range from <0.001-0.006 mg/day (Duggan and Corneliussen, 1972). The most likely source of pentachlorophenol contamination in many food products may be the exposure of the food to pentachlorophenol-treated wood materials such as storage containers (Dougherty, 1978).

Acute toxicity data indicated that salmonids are more sensitive to the toxic effects of pentachlorophenol than other fish species, with LC_{50} values of 34-128 µg/ℓ for salmonids and 60-600 µg/ℓ for other species. More recent data showed that carp larvae, bluegills, channel catfish and knifefish also had LC_{50} values <100 µg/ℓ (see Table 10). The most sensitive marine fishes were pinfish larvae, the goby, Gobius minutus, *and eggs and larvae of the flounder,* Pleuronectes platessa, *all with LC_{50} values <100 µg/ℓ (Adema and Vink, 1981). The most sensitive freshwater invertebrate species were the chironomid,* Chironomus gr. thummi *(Slooff, 1983) and the snail,* Lymnaea luteola *(Gupta et al., 1984). The most sensitive marine invertebrates were the Eastern oyster (Borthwick and Schimmel, 1978), larvae of the crustaceans,* Crangon crangon *and* Palaemon elegans *(VanDijk et al., 1977), and the copepod,* Pseudodiaptomus coronatus *(Hauch et al., 1980), all with LC_{50} values <200 µg/ℓ .*

In chronic toxicity tests, the lowest concentration reported to cause adverse effects was 1.8 µg/ℓ (NaPCP), which inhibited growth of sockeye salmon (Webb and Brett, 1973). The marine species tested displayed similar thresholds for chronic toxicity.

Both acute and chronic toxicity increased at lower pH, probably because a lower pH favors the un-ionized form of pentachlorophenol, which is taken up more readily and is therefore more toxic than ionized pentachlorophenol (Kobayashi and Kishino, 1980; Spehar et al., 1985).

Data concerning the effects of pentachlorophenol on aquatic plants were highly variable. Therefore, it was difficult to draw conclusions from these data.

Pentachlorophenol did not appear to bioaccumulate in aquatic organisms to very high concentrations. BCFs for pentachlorophenol were <1000 for most species tested. The highest BCF was 3830 for the polychaete, Lanice conchilega *(Ernst, 1979). Some species appear to have an inducible pentachlorophenol-detoxification mechanism, as evidenced in several experiments in which pentachlorophenol tissue levels peaked in 4–8 days and declined thereafter despite continued exposure (Pruitt et al., 1977; Trujillo et al., 1982). A study by Niimi and Cho (1983) indicated that uptake of waterborne pentachlorophenol from gills was much greater than uptake from food, indicating that bioconcentration of pentachlorophenol through the food chain is unlikely. Biomonitoring data of Lake Ontario fishes showed that similar pentachlorophenol levels were found in predators and forage species.*

Studies with experimental ecosystems have indicated that ecological effects may occur at pentachlorophenol levels as low as those causing chronic toxicity in sensitive species in single-species tests. The lowest concentration that caused adverse effects in these studies was 15.8 µg/ℓ , which caused a reduction in numbers of individuals and species in a marine benthic community (Tagatz et al., 1978).

Pentachlorophenol is readily absorbed from the gastrointestinal tract of rats, mice, monkeys and humans (Braun et al., 1977, 1978; Ahlborg et al., 1974; Braun and Sauerhoff, 1976). Peak plasma concentrations are reached within 12-24 hours after oral administration to monkeys (Braun and Sauerhoff, 1976), but 4-6 hours after oral administration to rats (Braun et al., 1977). After oral administration, the highest concentration of radioactivity was found in the liver and gastrointestinal tract of monkeys (Braun et al., 1977). In rats and mice, tetrachlorohydroquinone was identified in the urine (Jakobson and Yllner, 1971; Braun et al., 1977; Ahlborg et al., 1974) as well as unmetabolized pentachlorophenol and glucuronide-conjugated pentachlorophenol. Although Ahlborg et al. (1974) reported that oxidative dechlorination of pentachlorophenol occurs in humans, as evidenced by the presence of tetrachlorohydroquinone in the urine of workers occupationally exposed (probably by inhalation), analysis of human urine after ingestion of pentachlorophenol revealed the presence of conjugated pentachlorophenol and unmetabolized pentachlorophenol (Braun et al., 1978).

The primary route of excretion after oral administrtation of all species studied is in the urine (Braun et al., 1977, 1978; Ahlborg et al., 1974; Larsen et al., 1972; Braun and Sauerhoff, 1976). Although urinary excretion followed second-order kinetics in rats (Larsen et al., 1972;

Braun et al., 1977) except in females receiving a single high dose (100 mg/kg) of pentachlorophenol, urinary excretion of pentachlorophenol in humans and monkeys followed first-order kinetics (Braun and Sauerhoff, 1976; Braun et al., 1978). Enterohepatic circulation played an importation role in the pharmacokinetics of pentachlorophenol. The half-life of pentachlorophenol in the plasma is longer in female rats and monkeys than it is in male rats and monkeys (Braun et al., 1978; Braun and Sauerhoff, 1976).

Because many preparations of pentachlorophenol are contaminated with small but measurable amounts of highly toxic substances, such as dibenzodioxins, special attention must be paid to the composition of the pentachlorophenol solution tested. In studies where technical and purified pentachlorophenol have been evaluated (Schwetz et al., 1974; Goldstein et al., 1977; Kimbrough and Linder, 1978; Knudsen et al., 1974; Johnson et al., 1973; Kerkvliet et al., 1982), only the results of the experiments using purified pentachlorophenol were reported in detail. Oral exposure to pentachlorophenol was not carcinogenic in mice (BRL, 1968; Innes et al., 1969) or rats (Schwetz et al., 1977), regardless of the composition of the pentachlorophenol solution tested. Although there are a few studies that suggest pentachlorophenol may be mutagenic in B. subtilis *(Waters et al., 1982; Shirasu, 1976), in yeast,* Saccharomyces cerevisiae *(Fahrig et al., 1977) and in mice, as evidenced by the coat-color spot test (Fahrig et al., 1977), no evidence of mutagenicity was reported in* S. typhimurium *(Anderson et al., 1972; Simmon et al., 1977; Lemma and Ames, 1975; Moriya et al., 1983; Waters et al., 1982; Buselmaier et al., 1973) or in* E. coli *(Simmon et al., 1977; Fahrig, 1974; Moriya et al., 1983; Waters et al., 1982) with or without metabolic activation.*

Three teratogenicity/reproductive toxicity studies (Schwetz et al., 1974, 1977; Courtney et al., 1976) indicate that pentachlorophenol is fetotoxic in rats at oral dose levels $\geq$*5 mg/kg/day. At the highest dose tested (500 ppm) in a fourth teratogenicity/reproductive toxicity study (Exon and Koller, 1982), there was a statistically nonsignificant decrease in litter size. The lowest dose tested (5 mg/kg/day) by Schwetz et al. (1977) was the lowest dose at which any evidence of fetotoxicity, as indicated by delayed ossification, was observed. No adverse fetal or reproductive effects were reported at* $\leq$*3 mg/kg/day (Schwetz et al., 1977; Exon and Koller, 1982). In subchronic and chronic toxicity studies, adverse effects occurred primarily in the liver (Kerkvliet et al., 1982; Johnson et al., 1973; Knudsen et al., 1974; Goldstein et al., 1977; Kimbrough and Linder, 1978; Schwetz et al., 1977), the kidney (Johnson et al., 1973; Kimbrough and Linder, 1978; Schwetz et al.,*

1977) and the immune system (Kerkvliet et al., 1982). Knudsen et al. (1974) reported increased liver weights in female rats and centrilobular vacuolization in male rats exposed to diets containing ≥50 ppm commercial pentachlorophenol, which contained 282 ppm dioxins. In the remaining studies, increased liver weight (Johnson et al., 1973) and increased pigmentation of hepatocytes (Schwetz et al., 1977) were observed at oral doses of ≥10 mg/kg/day (~90%), and SGPT levels significantly increased in rats ingesting 30 mg/kg/day pentachlorophenol (~90%) for 2 years (Schwetz et al., 1977). Increased kidney weight unaccompanied by renal histopathology was reported in rats exposed to dietary concentration ≥20 ppm of pentachlorophenol (>99%) for 8 months (Kimbrough and Linder, 1978) and in rats ingesting 30 mg/kg/day (~90%) for 90 days (Johnson et al., 1973). Increased pigmentation of the renal tubular epithelial cells was reported in rats ingesting 10 or 30 mg/kg/day pentachlorophenol for 2 years (Schwetz et al., 1977). Although decreased immunocompetence was reported in mice exposed to dietary levels of 50 or 500 ppm of pentachlorophenol (>99%) for 34 weeks (Kerkvliet et al., 1982), the decrease was statistically significant only at the higher dose.

An ADI of 0.03 mg/kg/day or 2.1 mg/day for a 70 kg human was derived from the NOAEL of 3 mg/kg/day in rats in the chronic dietary study by Schwetz et al. (1977). An uncertainty factor of 100 was used. An RQ of 100 was derived based on the fetotoxic effects of pentachlorophenol in rats in the study by Schwetz et al. (1974). Based on guidelines for carcinogen risk assessment (U.S. EPA, 1984b) and inadequate evidence for animal carcinogenicity or absence of human cancer data, pentachlorophenol is classified as Group D, meaning that it is not classified as a human carcinogen.

INTRODUCTION

Structure and CAS Number. Pentachlorophenol is the common chemical name for 2,3,4,5,6-pentachlorophenol. It is commonly known by the synonyms PCP or penta, or by the trade names Chlorophen, Dowicide 7 or Santophen 20. The structure, molecular weight, empirical formula and CAS Registry number for pentachlorophenol are as follows:

Molecular weight: 266.34
Empirical formula: C_6HCl_5O
CAS Registry number: 87-86-5

TABLE 1
Bioconcentration of Pentachlorophenol in Aquatic Organisms

Organism	BCF	Comments	Reference
Carp	300-400	3-day static-tank test	Gluth et al., 1985
Fish (not specified)	200	Test parameters not given	Branson, 1980
Clams	100-1000	Determined from clams and water taken from a natural water body which had been receiving pentachlorophenol effluents	Butte et al., 1985
Rainbow trout	5360 39-615	Monitoring of bile after 24-hour exposure 24-hour static test; low BCF for muscle, high BCF for liver	Statham et al., 1976; Carlson et al., 1979
Fathead minnow	770	32-day exposure in a flowthrough aquaria	Veith et al., 1979a
Lanice conchilega S. troglodytes	2600-8500 70-180	Both organism types taken from the Weser estuary in Germany with BCF levels determined for the natural water containing 0.04 ng/ℓ pentachlorophenol	Ernst and Weber, 1978
Killfish (Fundulus similus)	10-64	4- to 168-hour exposure in aquaria; depuration half-life of 4.7 days, determined following pentachlorophenol exposure	Trujillo et al., 1982
Alga (Chlorella fusca)	1250	24-hour static test	Geyer et al., 1981

TABLE 1 (continued)

Organism	BCF	Comments	Reference
Fish (*Gambusia*)	297	24-hour model aquatic ecosystem study	Lu and Metcalf, 1975
Snail (*Physa*)	121		
Mosquito (*Culex*)	17		
Water flea (*Daphnia*)	165		
Alga (*Oedogonium*)	2		
Golden orfe fish	350-1100	Static test; 350 BCF for 24 hours; 1100 BCF for 3-day period	Korte et al., 1978
Goldfish	180	12-hour static test	Kobayashi et al., 1979
Eastern oyster	64	NC	Schimmel and Garnas, 1981
Fish	10,000	Determined from fish and water samples taken from the Bay of Quinte, Lake Ontario	Fox and Joshi, 1984

NC = no comment

Physical and Chemical Properties: Pure pentachlorophenol is a white monoclinic crystalline solid with a phenolic odor; the technical grade material is dark grey to brown (Verschueren, 1983; Martin and Worthing, 1977). Selected physical properties are listed below:

Melting point	190°C	Freiter, 1978
Boiling point:	310°C (decomposes)	Verschueren, 1983
Specific gravity:	1.978	Verschueren, 1983
Water solubility:		
at 0°C	5 ppm	Verschueren, 1983
at 20°C	14 ppm	U.S. EPA, 1981a
at 30°C	20 ppm	Martin and Worthing, 1977
at 50°C	35 ppm	Verschueren, 1983
Vapor pressure (mm Hg):		
at 0°C	1.7×10^{-5}	Crosby et al., 1981
at 20°C	1.7×10^{-4}	Crosby et al., 1981
at 20°C	1.1×10^{-4}	U.S. EPA, 1981a
at 20°C	1.4×10^{-4}	Lyman et al., 1982
Log K_{ow}	5.12	Hansch and Leo, 1985
pK_a (20-25°C):	4.70	Cessna and Grover, 1978
	4.74	Drahonovsky and Vacek, 1971
Henry's Law constant:	3.4×10^{-6} atm•m^3/mol (calculated)	Lyman et al., 1982

Technical grade pentachlorophenol typically contains 85-90% pentachlorophenol, 6-12% tetrachlorophenol and other polychlorophenols, and traces to several thousand ppm of polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans (Crosby et al., 1981; IARC, 1979).

As with any phenol, the hydroxyl group of pentachlorophenol will undergo nucleophilic reactions; it forms esters with organic acids, ethers with alkylating agents and salts with alkaline metals (Crosby et al., 1981; IARC, 1979). The ring chlorines are resistant to nucleophilic displacement under normal conditions.

Production Data. Pentachlorophenol was introduced as a wood preservative in 1936 (Martin and Worthing, 1977); however, its commercial production in the United States was not reported by the U.S. Tariff Commission until 1950 (IARC, 1979). The most recent production figure available is 46.8 million pounds in 1980. The manufacturers of pentachlorophenol in 1980 were cited as follows (CMR, 1980):

	Annual Capacity (million lbs)
Dow Chemical (Midland, MI)	30
Reichhold Chemical (Tacoma, WA)	18
Vulcan Chemical (Wichita, KA)	20

Reichhold and and Vulcan, however, are the only current manufacturers (SRI, 1985).

Pentachlorophenol is manufactured by the chlorination of phenol in the presence of various catalysts (Freiter, 1978).

Use Data. Pentachlorophenol is used as an industrial wood preserative for utility poles, crossarms, fence posts, etc. (79%): for NaPCP (12%); and miscellaneous, including mill uses, consumer wood preserving formulations and herbicide intermediate (9%) (CMR, 1980). As a wood preservative, pentachlorophenol acts as both a fungicide and insecticide (Freiter, 1978). The miscellaneous mill uses primarily involve the application of pentachlorophenol as a slime reducer in paper and pulp milling and may constitute ~6% of the total annual consumption of pentachlorophenol (Crosby et al., 1981). NaPCP is also used as an antifungal and antibacterial agent (Freiter, 1978). Pentachlorophenol also is used as a general herbicide (Martin and Worthing, 1977).

ENVIRONMENTAL FATE AND TRANSPORT PROCESSES

Water. *Hydrolysis.* Based on the chemical structure activity relationships of pentachlorophenol, hydrolysis is not expected to be environmentally significant (Callahan et al., 1979a; Mabey et al., 1981).

Oxidation. Experimental data regarding the oxidation of pentachlorophenol were not located; however, highly chlorinated organic compounds are usually resistant to oxidation at ambient water temperatures. Therefore, oxidation of pentachlorophenol is not expected to be environmentally important (Callahan et al., 1979). Mabey et al. (1981) theoretically calculated the rate constants for the reactions between pentachlorophenol and singlet oxygen ($^1O_2$) and pentachlorophenol and peroxy radical ($RO_2\bullet$) at 25°C to be $<7x10^3$ and $1x10^5$ M^{-1} hr^{-1}, respectively. These rate constants indicate that oxidation may not be important.

Photolysis. A substantial amount of experimental data indicates that photolysis is a major removal process of pentachlorophenol in water. Aqueous solutions of pentachlorophenol absorb radiation quite strongly within the sunlight region of the spectrum, with the absorption maximum occurring at 320 nm (ECETOC, 1984; Wong and Crosby, 1981).

Pentachlorophenol is a moderately acidic substance (pK_a of 4.74) (Drahonovsky and Vacek, 1971), and will exist primarily as an anion in environmental surface waters (Callahan et al., 1979a). Wong and Crosby (1981) found that the rate of photolysis of the anion is much faster than that of the undissociated compound. In aqueous

solutions of pH 7.3 and 3.3 exposed to laboratory UV light, the pentachlorophenol half-lives were 3.5 and ~100 hours, respectively. Exposure of the pH 7.3 solution to July sunlight at Davis, CA, resulted in 50% degradation in 48 hours and total disappearance within 10 days. Wong and Crosby (1981) also identified a variety of products resulting from the photodegradation of pentachlorophenol or its metabolites; these degradation products included 2,3,4,6- and 2,3,5,6-tetrachlorophenol, tetrachlorocatechol, tetrachlororesorcinol, tetrachlorohydroquinone, chloranil, hydroxyqinonrd and 2,3-dichloromaleic acid.

ECETOC (1984) measured the rate of photolysis of a dilute aqueous solution in a specially designed laboratory apparatus and found pentachlorophenol to photodegrade rapidly. The experimental results were used to theoretically calculate a yearly-averaged, photolytic rate constant (at surface conditions at a latitude of 40-50° N) of 0.00195 be 2.1 hr^{-1} in midday sunlight, which corresponds to a half-life of 0.33 hours. It should (1984) reported the direct aqueous photolytic rate constant for pentachlorophenol to be 2.1 hr^{-1} at midday sunlight, which corresponds to a half-life of 0.33 hours. It should be noted that the rapid photolysis rates reported by ECETOC (1984) and Zepp et al. (1984) are for surface photolysis only; as water depth increases, the photolysis rate decreases. Therefore, it is possible for photolysis to be the dominant removal mechanism in shallow bodies of water and only a secondary or minor removal mechanism in deep lakes or the ocean.

Pignatello et al. (1983) examined the photolysis rate (as well as other fate processes) of pentachlorophenol in outdoor manmade channels at a U.S. EPA field station on the Mississippi River near Monticello, MN. The photolytic half-lives at various water depths were determined as follows:

Water Depth (cm)	3-Hour Average Half-Life (10 a.m. to 1 p.m.)	24-Hour Average Half-Life
0.5	0.70	3.02
3.8	2.40	4.20
7.2	3.73	16.9
13.8	9.63	43.1

This clearly demonstrates that photolysis decreases significantly as water depth increases.

Crossland and Wolff (1985) studied the fate of pentachlorophenol in outdoor ponds to which the compound was added. Using the SOLAR mathematical model, the authors theoretically calculated that the photolysis rate constant would range between 0.23 and 0.46 day^{-1} in the ponds, which was similar to the observed rate of 0.15-0.34 day^{-1}. It was concluded that phototransformation of pentachlorophenol could be an important removal mechanism in northern temperate climates.

Pierce and Victor (1978) monitored the fate of pentachlorophenol and its degradation products that had been accidentally released into a freshwater lake from a settling pond that held wood-treating wastes. The origin of two of the major degradation products (2,3,5,6- and 2,3,4,5-tetrachlorophenol) was attributed to photolytic dechlorination of pentachlorophenol while it was still being held in the settling pond. Hydrocarbon wastes were also detected.

Kotzias et al. (1982) compared the photolysis rate of pentachlorophenol in pure water to water containing nitrite. Under laboratory conditions, the half-life in pure water was 0.46 hours, while the half-life in nitrite was only 0.2 hours, suggesting that photolysis in natural water may be accelerated by the presence of specific, naturally occurring materials.

Microbial Degradation. Banerjee et al. (1984) examined the biodegradation of pentachlorophenol in two natural river waters under laboratory conditions. In Seneca River water, pentachlorophenol biodegraded significantly after a lag period of 15 days; the Seneca River was noted to receive both domestic and industrial effluents and, therefore, was likely to contain organisms adapted to various organic compounds. In the relatively unpolluted Jamesville River water, however, pentachlorophenol did not biodegrade. Baker et al. (1980) tested the aerobic degradation of pentachlorophenol in a natural stream water for 40 days and found virtually no degradation in either sterile or nonsterile samples.

Tabak et al. (1981) used a static culture flask-screening procedure to test the biodegradation and acclimation periods of the U.S. EPA priority pollutants. Pentachlorophenol was found to undergo significant biodegradation with gradual adaptation; 99-100% BOD rates were determined for concentrations of 5 and 10 ppm and 21 and 28 days' incubation, respectively.

Rott et al. (1982) reported the results of two laboratory biodegradation tests with respect to pentachlorophenol. In the GSF test (5 days incubation, 0.05 ppm), the biodegradation rate was only 0.1% of theoretical CO_2. The modified OECD test (28 days incubation, 5-40 ppm) also showed slow biodegradation with a DOC rate of 5%.

Properly acclimated activated sludges and mixed bacterial cultures have been shown to significantly biodegrade pentachlorophenol in water (Ingols et al., 1966; Hickman and Novak, 1984; Madden and Tittlebaum, 1983; Moos et al., 1983; Klecka and Maier, 1985). The U.S. EPA (1981a) reported that 15 full-scale wastewater treatment plants and 1 pilot plant using activated sludges remove 67 to $\geq$99% of influent pentachlorophenol. Korte and Klein (1982) found a slow biodegradation rate (0.5% of theoretical CO_2 in 5 days) in an activated sludge system test using unacclimated sludge.

A variety of fungal and bacterial isolates that may be present in natural waters are capable of biodegrading pentachlorophenol (Cserjesi and Johnson, 1972; Cook et al., 1980; Edgehill, 1982; Karns et al., 1983; Kwasniewska, 1981; Duncan and Deverall, 1964). The fungus *Trichoderma virgatum* has been reported to produce the metabolite pentachloroanisole by the methylation of pentachlorophenol (Cserjesi and Johnson, 1972).

Pignatello et al. (1983) monitored most of the aquatic fate mechanisms in outdoor manmade channels: Microbial degradation of pentachlorophenol was found to become significant 3 weeks after initial dosing, and eventually became the dominant removal mechanism. Bourquin (1984) tested the biodegradation of pentachlorophenol in three sea waters and one river water using an eco-core system test; degradation in sterile controls was significantly slower than in nonsterilized waters, indicating microbial removal.

Volatilization. Sugiura et al. (1984) experimentally measured the overall K_L for pentachlorophenol from water to be 0.023 cm/hr at 20.7°C and wind velocity of 1.7 m/sec. The volatilization half-life can be calculated by (Lyman et al., 1982):

$$t_{1/2} = \frac{0.6931 \text{ x (mean depth of water body)}}{K_L}$$

In a body of water 1 m deep, the half-life is calculated to be ~125 days.

Moos et al. (1983) monitored pentachlorophenol volatilization losses from continuously stirred biodegradation reactors over a 2-week period to determine the importance of evaporation. It was determined that the maximum loss of pentachlorophenol from volatilization was only 0.037% of the initial concentration. Pignatello et al. (1983) found minor losses of pentachlorophenol that were due to volatilization from outdoor aquatic channels; volatilization loss in the form of vapor and aerosol was considered to be <0.006%.

These experimental data indicate that volatilization is not an important environmental fate mechanism in relation to photolysis or biodegradation.

Adsorption. Schellenberg et al. (1984) conducted laboratory experiments to study the sorption of chlorinated phenols by sediments and aquifer materials. It was demonstrated that sorption of the conjugate bases (phenolates) occurred as well as sorption of the nondissociated phenols. Pentachlorophenol is present predominately in the phenolate form at ambient pH values, but sorption in this form can occur with the degree of sorption dependent on the ionic strength in the aqueous phase (Schellenberg et al., 1984). Experimental studies by DeLaune et al. (1983) suggested that pentachlorophenol is more tightly bound by oxidized sediment solids than by reduced sediment solids; surface sediment and colloidal suspended particulates tend to be oxidized.

Pierce and Victor (1978) reported on the fate of accidental spills of pentachlorophenol into a freshwater lake. Sediments and leaf litter retained high concentrations of pentachlorophenol throughout the monitoring period (8 months) and were considered, along with leaching, as sources of continuing pollution into the lake.

The above data suggest that the sorption and desorption of pentachlorophenol on organic aquatic particulate matter and sediments may have some environmental significance.

Bioaccumulation. A substantial number of studies have measured the bioconcentration of pentachlorophenol in aquatic organisms. Table 1 lists the measured BCFs determined

by various studies and indicates that pentachlorophenol will bioaccumulate in various aquatic organisms. This is consistent with the widespread detection of pentachlorophenol in fish and other organisms taken from ambient bodies of water (Murray et al., 1980, 1981; Kuehl et al., 1980, 1983; Gossett et al., 1983; Veith et al., 1979a; Fox and Joshi, 1984; Niimi and Cho, 1983; Butte et al., 1985; Zitko et al., 1974; Pierce and Victor, 1978; Ernst and Weber, 1978; IARC, 1979; Eder and Weber, 1980; New Brunswick Res. Prod. Council, 1978; Young et al., 1983; Paasivirta et al., 1980; Renberg et al., 1983; Ray et al., 1983a,b).

Air. *Chemical Removal Processes.* The UV spectrum for nondissociated pentachlorophenol shows some absorption >290 nm (Sadtler, 1966), indicating a potential for atmospheric photolysis. Wong and Crosby (1981) found that the ionized form of pentachlorophenol, which exists in ambient bodies of water, photolyzes much faster in aqueous solution than the nondissociated form; however, the nondissociated form did photolyze at an environmentally significant rate, which indicates that photolysis of the nondissociated form in the atmosphere may be significant. Korte and Parlar (1977) found that pentachlorophenol adsorbed to solid particles photomineralized significantly in laboratory tests designed to predict photodecomposition in the atmosphere. Therefore, it is possible that direct photolysis is an important environmental sink for pentachlorophenol present in the atmosphere.

Data regarding the reaction of pentachlorophenol with hydroxyl radicals or with ozone in the air could not be located in the available literature as cited in the Appendix. It is likely that OH radicals will react with pentachlorophenol; however, the rate of reaction cannot be predicted without experimental data.

Physical Removal Processes. Pentachlorophenol has been detected in snow and rain (Paasivirta et al., 1985; Bevenue et al., 1972), suggesting that removal from the atmosphere by the wash-out effect of rain or by dissolution into clouds with subsequent precipitation can occur.

Soil. *Microbial Degradation.* The significant amount of available data indicate that microbial degradation is an important, and potentially dominant, removal mechanism for pentachlorophenol in soil. Baker et al. (1980) examined the aerobic degradation of pentachlorophenol remained in either the free or conjugated form. The identified degradation products included tetrachlorophenol, trichlorophenol, and tri-, tetra- and pentachloroanisole.

Edgehill and Finn (1983) examined the microbial degradation of pentachlorophenol in a moist garden. The half-life in nonsterile soil was ~12 days with only a 30% decomposition rate over the same time period in sterile soil. Addition of *Arthrobacter* cells acclimated to pentachlorophenol reduced the half-life to only 1 day. Alexander and Aleem (1961) used laboratory tests to determine that complete microbial decomposition of pentachlorophenol in Dunkirk soil required >72 days. Kirsch and Etzel (1973) have shown that microorganisms obtained from soil previously exposed to pentachlorophenol are capable of biodegrading pentachlorophenol.

Various bacteria species isolated from soils have also been shown to be capable of biodegrading pentachlorophenol (Watanabe, 1973; Trevors, 1982; Chu and Kirsch, 1973). Ahlborg and Thunberg (1980) reported that degradation of pentachlorophenol in submerged soils does not occur in sterilized soils, suggesting that under submerged conditions, microbial decomposition is the only significant chemical removal mechanism.

Photolysis. Gaeb et al. (1975) demonstrated that pentachlorophenol adsorbs to silica gel and undergoes significant photomineralization when exposed to laboratory UV radiation >290 nm. Lamparski et al. (1980) exposed pentachlorophenol-treated wood to sunlight and found that the pentachlorophenol was susceptible to photolysis, yielding octachlorodibenzo-p-dioxin as a photodegradation product. These data indicate that pentachlorophenol on soil surfaces exposed to sunlight is susceptible to photodegradation.

Adsorption. Using the soil TLC method, Helling et al. (1974) and Dragun and Helling (1981) determined the soil R_f (refractive index) value for pentachlorophenol in Hagerstown silty clay loam to be 0.40-0.42; this R_f value classified pentachlorophenol as having intermediate mobility in soil. Kenaga (1980) reported an experimental K_{oc} value of 900 for pentachlorophenol, which indicates low soil mobility.

Pentachlorophenol is relatively immobile in acidic Hawaiian soil, but mobility increases as pH increases (Sanborn et al., 1977). Callahan et al. (1979) also reported the increased adsorption in acidic soil systems as compared with neutral soils. Pentachlorophenol exists primarily as the phenolate ion in neutral aqueous solutions; however, the nondissociated form increases as the solution becomes more acidic. In general, the nondissociated form is much more tightly sorbed to sediments in aqueous systems than the ionized form, although the ionized form can experience significant sorption under proper conditions (Schellenberg et al., 1984). Therefore, pentachlorophenol appears to be susceptible to leaching in neutral or alkaline soils, but adsorption is expected to increase in more acidic soils.

Volatilization. Although experimental soil volatilization data suggest that pentachlorophenol evaporation from soil surfaces may have some importance (Kilzer et al., 1979), volatilization is not expected to be generally important. Soil volatility rates are dependent on factors such as moisture content, soil type, temperature and weather.

Persistence and Degradation. Pentachlorophenol decomposes more rapidly in soils of high organic content than in soils of low organic content, and more rapidly when moisture content is high and temperature is conducive to microbial activity. Half-lives are usually ~2–4 weeks (Crosby et al., 1981).

Weiss et al. (1982) examined the degradation of pentachlorophenol added to flooded rice soil in a plant growth chamber. After one vegetation season, only 1.12% of applied ceptible to adsorption in some soils (Schellenberg et al., 1984). In laboratory soils, pentachlorophenol decomposes faster in soils of high organic content as compared to low organic content, and faster when moisture content is high and the temperature is

conducive to microbial activity. Half-lives are usually ~2–4 weeks (Crosby et al., 1981).

Ide et al. (1972) found that pentachlorophenol applied as an herbicide to rice fields decomposed within a few weeks after application, with microbial degradation being the dominant decomposition process. The stable decomposition products identified in the paddy soil included 2,3,4,5-, 2,3,5,6- and 2,3,4,6-tetrachlorophenol; 2,4,5- and 2,3,5-trichlorophenol; 3,4- and 3,5-dichlorophenol; and 3-chlorophenol.

Suzuki and Nose (1970) observed that 90% of pentachlorophenol applied to a farm soil at 100 ppm disappeared in 10 days, while only 10% disappeared when applied at 1000 ppm. Murthy et al. (1979) examined both the aerobic and anaerobic degradation of pentachlorophenol in the laboratory using Hagerstown silty clay loam soil. Pentachlorophenol was a degradation product of both aerobic and anaerobic conditions with the metabolites, 2,3,5,6- and 2,3,4,5- tetrachlorophenol and 2,3,6-trichlorophenol, present under anaerobic conditions.

Middleton (1984) described the case history of a wood-treating site where the soil became contaminated with pentachlorophenol at a concentration of 700 ppm. The site was prepared for *in situ* biodegradation by fertilizer application, tilling and moisture control. Within a 140-day period, the pentachlorophenol concentration of the soil dropped to 140 ppm.

Summary. Photolysis and microbial degradation are the important chemical removal mechanisms for pentachlorophenol in water. In surface waters, pentachlorophenol photolyzes rapidly (ECETOC, 1984; Wong and Crosby, 1981; Zepp et al., 1984); however, the photolytic rate decreases as the depth in water increases (Pignatello et al., 1983). Pentachlorophenol is readily biodegradable in the presence of acclimated microorganisms; however, biodegradation in natural waters requires the presence of microbes that can become acclimated. A natural river water that had been receiving domestic and industrial effluents significantly biodegraded pentachlorophenol after a 15-day lag period, while an unpolluted natural river water was unable to biodegrade the compound (Banerjee et al., 1984). Even though pentachlorophenol is in the ionized form in natural waters, sorption to organic particulate matter and sediments can occur (Schellenberg et al., 1984), with desorption contributing as a continuing source of pollution in a contaminated environment (Pierce and Victor, 1978). Experimentally determined BCFs have shown that pentachlorophenol can significantly accumulate in aquatic organisms (Gluth et al., 1985; Butte et al., 1985; Statham et al., 1976; Veith et al., 1979a,b; Ernst and Weber, 1978), which is consistent with its widespread detection in fish and other organisms.

Direct photolysis may be an important environmental sink for pentachlorophenol present in the atmosphere. The detection of pentachlorophenol in snow and rain water (Paasivirta et al., 1985; Bevenue et al., 1972) suggests that removal from air by dissolution is possible.

Soil degradation studies indicate that pentachlorophenol is biodegradable, and microbial decomposition is an important and potentially dominant removal mechanism in soil (Baker et al., 1980; Baker and Mayfield, 1980; Edgehill and Finn, 1983; Kirsch and Etzel, 1973; Ahlborg and Thunberg, 1980). The degree to which pentachlorophenol leaches in soil is dependent on the type of soil. In soils of neutral pH, leaching may be significant, but in acidic soils, adsorption to soil generally increases (Callahan et al., 1979a; Sanborn et al., 1977). The ionized form of pentachlorophenol may be sus- pentachlorophenol in a sediment sample for a 30-day period; degradation of nonsterile sediment was 24-29%, while degradation in sterile sediment was only 12-18%. Baker and Mayfield (1980) conducted similar laboratory tests using a clay loam soil; after 160 days' incubation, 80% of the pentachlorophenol was degraded in the nonsterile soil and only 20% degraded in the sterile soil.

EXPOSURE

Water. Pentachlorophenol has been detected in surface waters, groundwater, drinking water and effluents from sewage and industrial sources (Table 2). In addition, pentachlorophenol has been detected in widespread aquatic sediment samples (Eder and Weber, 1980; Watanabe et al., 1985; Xie, 1983; Pierce et al., 1980; Ray et al., 1983a,b; Wegman and Vandenbroek, 1983).

Cole et al. (1984) reported preliminary findings of the U.S. EPA's National Urban Runoff Program, which has monitored urban storm water runoff. Pentachlorophenol has been detected in 15% of the samples at concentrations ranging from 1-115 μg/ℓ. Pentachlorophenol was detected in Long Island, NY; Austin, TX; Little Rock, AR; Bellevue, WA and Eugene, OR.

The U.S. EPA conducted the National Organic Monitoring Survey to determine the frequency of specific contaminants in the drinking water of 113 public water supplies (Brass et al., 1977). During phases I and II of the survey, pentachlorophenol was identified in 91/118 analyses with mean concentrations of 0.008-0.07 μg/ℓ (Mello, 1978). The gross analysis of data from the U.S. EPA STORET Data Base is listed in Table 3.

There are at least several sources by which pentachlorophenol can be released to water. A primary source is likely to be the use of pentachlorophenol in wood preservation and the effluents associated with this industry. The use of pentachlorophenol as an herbicide directly exposes the environment to the compound. Murthy and Kaufman (1978) identified pentachlorophenol as a degradation product of the pentachloronitroben- zene; this may be a potential indirect source of pentachlorophenol exposure to water.

Inhalation. Pentachlorophenol can be emitted to the atmosphere by evaporation from treated wood or water surfaces, by releases from cooling towers using pentachlorophenol as a biocide or by incineration of treated woods (Skow et al., 1980; Crosby et al., 1981).

TABLE 2
Water Monitoring of Pentachlorophenol

Location	Water Source	Concentration (ppm)	Reference
Finland	lake waters	0-0.006	Paasivirta et al., 1985
Netherlands	surface waters	0.01 (maximum)	Wegman and Hofstee, 1979
West Germany	river water	0.0005 (maximum)	Weber and Ernst, 1978
United States (not specified)	sewage effluent	NR	U.S. EPA, 1975
United States (not specified)	industrial wastewater	0-8.3	U.S. EPA, 1981a
Sweden	river waters	0.091 (maximum)	Renberg et al., 1983
Japan	river waters	0-0.0003	Matsumoto, 1982
Dade County, FL	drinking water	0-0.0003	Morgade et al., 1980
Los Angeles, CA	municipal wastewater effluent	0-0.003	Young et al., 1983
Switzerland	river water	0-0.00008	Schwarzenbach et al., 1983
Texas	marine water	0.000004-0.000011	Murray et al., 1981
Germany	river water	0.0004 (maximum)	Malle, 1984
United States (not specified)	sewage effluent	0.00026 (average)	Callahan et al., 1979b
Oregon	sewage effluent	0.0001-0.00044	Buhler et al., 1973

TABLE 2 (continued)

Location	Water Source	Concentration (ppm)	Reference
Wisconsin	wastewaters	0.0001-0.0040	Peterman et al., 1980
Texas (creosote waste site)	groundwaters	0-0.0584	Bedient et al., 1984
NR	drinking water	NR	Kool et al., 1982
Finland	surface waters	0-0.0003	Valo et al., 1984
Texas	estuary waters	0.00068 (maximum)	Ray et al., 1983a,b
United States (not specified)	industrial and sewage effluents	NR	Shackelford and Keith, 1976
Switzerland	pulp mill effluent	0.0007-0.002	Leuenberger et al., 1985
Oroville, CA	drinking water	0.227 (average)	Wong and Crosby, 1981
United States (not specified)	groundwater	0.040	Stover and Kincannon, 1983
Finland	pulp effluents	0.0022 (maximum)	Salkinoja-Salonen et al., 1984
Germany	river water	0.00000004 (average)	Ernst and Weber, 1978
Lake Ontario	lake water	0.00011 (maximum)	Fox and Joshi, 1984

NR = Not reported

TABLE 3
Gross Analysis of Pentachlorophenol in Water, Sediment and Aquatic Organisms[a]

Concentration	Whole Water[b] (ppm)	Water Only[c] (ppm)	Sediment[d] (mg/kg dry weight)	Aquatic Organism Tissue[e] (mg/kg wet weight)
Maximum	203	0.010	570	2000
Minimum	0	0.007	0	0
Mean	0.138	0.0099	2	3

[a]Source: U.S. EPA, 1985a
[b]N = 7369
[c]N = 194
[d]N = 1511
[e]N = 2226

Pentachlorophenol has been detected in snow and rain water (Paasivirta et al., 1985; Bevenue et al., 1972), evidence of its presence in ambient atmospheres. Cautreels et al. (1977) detected pentachlorophenol in the air of Antwerp, Belgium (5.7-7.8 ng/m^3) and Chacaltaya, Bolivia (0.25-0.93 ng/m_3). The indoor air of dwellings treated with pentachlorophenol to control wood boring insects has been found to contain pentachlorophenol (Dobbs and Williams, 1983). Pentachlorophenol has been found in occupational atmospheres (IARC, 1979; Crosby et al., 1981) and identified in emissions from hazardous waste incineration (Oberg et al., 1985).

Food. The U.S. Food and Drug Administration initiated the Total Diet Program (Market Basket Study) in 1964 to monitor ready-to-eat foods for pesticide residues. The results of these monitoring data with respect to pentachlorophenol are listed in Table 4 (adult diets) and Table 5 (infant and toddler foods).

Duggan et al. (1983) compiled the results (for fiscal years 1970-1976) of three Federal programs designed to monitor chemical residues in foods; these programs include the Total Diet Program, the U.S. FDA monitoring program for raw agricultural commodities, both domestic and imported, and the U.S. Department of Agricultural National Residue Program, which examined meats and poultry products. Pentachlorophenol was found infrequently in dairy products, fruits, leaf and stem vegetables, vine and ear vegetables, root vegetables, processed vegetable products and miscellaneous processed foods. Dugan and Corneliussen (1972) estimated that the average dietary intake of pentachlorophenol in the United States during the years 1965-1969 ranged from <0.001-0.006 mg/day.

TABLE 4
Pentachlorophenol Residues in Total Diet Samples*

Dates of Food Collection	Total Number of Food Composites Analyzed	Number of Composites with Residues	Number of Positive Composites with Residues Reported as Trace	Concentration Range (ppm)	Food Classes Containing Residues
June 1964-April 1965	216	3	1	0.02-0.03	meats, grain and cereals
June 1965-April 1966	432	10	5	0.036-0.31	dairy products, grain and cereals, oils and fats, beverages
June 1966-April 1967	360	11	2	0.007-0.043	dairy products, meats, grain and cereals, potatoes, sugars and adjuncts, beverages
June 1967-April 1968	360	7	1	0.009-0.14	dairy products, meats, grain and cereal, potatoes
June 1968-April 1969	360	10	5	0.02-0.04	oils and fats, garden fruits, sugars and adjuncts, grains and cereals, dairy products, meats and fish
June 1969-April 1970	NR	NR	NR	NR	NR
June 1970-April 1971	360	2	1	trace to 0.011	dairy products, sugar and adjuncts
June 1971-July 1972	NR	NR	NR	NR	NR
August 1972-July 1973	360	9	1	0.01-0.02	dairy products, legume vegetables, sugar and adjuncts
August 1973-July 1974	360	10	0	0.01-0.033	dairy products, grain and cereals, root vegetables, sugar and adjuncts
August 1974-July 1975	360	13	0	0.01-0.04	dairy products, grain and cereals, leafy and root vegetables, garden fruits, sugar and adjuncts
August 1975-July 1976	240	2	0	0.026-0.06	sugar and adjuncts, beverages
August 1976-September 1977	360	1	0	0.016	grain and cereals
October 1977-September 1978	NR	NR	NR	NR	NR

*Sources: Duggan et al., 1966, 1967; Martin and Duggan, 1968; Corneliussen, 1969, 1970, 1972; Manske and Corneliussen, 1974; Manske and Johnson, 1975, 1977; Johnson and Manske, 1976, 1977; Johnson et al., 1981a, 1984a; Podrebarac, 1984a

NR = Not reported

TABLE 5
Pentachlorophenol Residues in Infant and Toddler Total Diet Samples*

Dates of Food Collection	Total Number of Food Composites Analyzed	Number of Composites with Residues	Number of Positive Composites with Residues Reported as Trace	Concentration Range (ppm)	Food Classes Containing Residues
August 1974-July 1975	220	2	0	0.01-0.09	vegetables, sugar and adjuncts
August 1975-July 1976	220	10	9	trace to 0.07	whole milk, other dairy, grain and cereals, fruit and fruit juices, vegetables, sugar and adjuncts
August 1976-September 1977	220	1	0	0.035	sugar and adjuncts
October 1977-September 1978	NR	NR	NR	NR	NR

*Sources: Johnson et al., 1979, 1981b, 1984b; Podrebarac, 1984b
NR = Not reported

Dougherty (1978) reported that a spot survey of selected items resulted in the detection of pentachlorophenol residues in powdered milk, soft drinks, bread, candy bars, cereal, noodles, rice, sugar and wheat; concentrations ranged from 1 ppb to 0.1 ppm. As noted previously, pentachlorophenol has been detected frequently in fish and other aquatic organisms that may be potential food products. Gebefügi and Korte (1983) detected pentachlorophenol in human breast milk.

Dougherty (1978) also suggested that the most likely source of pentachlorophenol contamination in many foods is the exposure of the food to wood treated with pentachlorophenol. The presence of pentachlorophenol residues in grain and sugar products is consistent with the storage of these products in pentachlorophenol-treated wooden storage containers (Dougherty, 1978).

Dermal. Pertinent data regarding the monitoring of dermal exposure to pentachlorophenol could not be located in the available literature as cited in the Appendix; however, workers involved with the pentachlorophenol treatment of wood products or its use as a pesticide are likely to be exposed, along with consumers coming in contact with pentachlorophenol-treated products.

Summary. Monitoring studies have confirmed the widespread occurrence of pentachlorophenol in surface waters, groundwater, drinking water and industrial effluents (see Table 2). The U.S. EPA's National Urban Runoff Program and National Organic Monitoring Survey reported frequent detections in storm water runoff and public water supplies (Cole et al., 1984; Mello, 1978). Primary sources by which pentachlorophenol may be emitted to environmental waters may be through its use in wood preservation and the associated effluents and its pesticidal applications. Pentachlorophenol can be emitted to the atmosphere by evaporation from treated wood or water surfaces, by releases from cooling towers using pentachlorophenol biocides or by incineration of treated wood (Skow et al., 1980; Crosby et al., 1981). Pentachlorophenol has been detected in ambient atmospheres (Cautreels et al., 1977), in snow and rain water (Paasivirta et al., 1985; Bevenue et al., 1972) and in emissions from hazardous waste incineration (Oberg et al., 1985). The U.S. Food and Drug Administration's Total Diet Study (conducted between 1964 and 1977) found pentachlorophenol residues in 91/4428 ready-to-eat food composites (see Tables 4 and 5). The average American dietary intake of pentachlorophenol during 1965-1969 was estimated to range from <0.001-0.006 mg/day (Duggan and Corneliussen, 1972). The most likely source of pentachlorophenol contamination in many food products may be the exposure of the food to pentachlorophenol-treated wood materials such as storage containers (Dougherty, 1978).

PHARMACOKINETICS

Absorption. Results of studies using rats, mice, monkeys and humans (Braun et al., 1977, 1978; Ahlborg et al., 1974; Braun and Sauerhoff, 1976) indicate that pentachlorophenol is readily absorbed by the gastrointestinal tract, as evidenced by urinary

TABLE 6
Distribution Activity After Intraperitoneal Injection of [^{14}C]Pentachlorophenol in the Female NMRI Mice[a,b]

Dose (mg/kg)	Percent of Dose									
	Liver	Kidney	Lung	Heart	Brain	Stomach[c]	Gall Bladder[c]	Intestines[c]	Residue	Total
14.8	2.2 [9]	0.09 [2]	0	0.03 [2]	0.01 [0.1]	NR	NR	NR	5.0	98.1[d]
18.2	2.5 [10]	0.5 [8]	NR	NR	NR	2.1 [21]	0.08 [60]	4.8 [8]	0.6	87.6[d]
37.2	3.1 [26]	0.04 [1]	NR	0	NR	0.5 [10]	0.06 [90]	2.2 [8]	2.0	84.8[d]
35.2	1.2 [10]	0.04 [1]	0.02 [1]	0.03 [3]	0.02 [0.5]	NR	NR	NR	1.4	93.9[e]
36.8	0.4 [3]	0.01 [0.3]	0	0.01 [1]	0	NR	NR	NR	0.1	92.4[f]

[a]Source: Jakobson and Yllner, 1971
[b][ ]Activity equivalent to μg pentachlorophenol/g organ weight
[c]With contents
[d]Killed after 96 hours
[e]Killed after 7 days
[f]Killed after 30 days, urine and feces analyzed for the first week
NR = Not reported

excretion data. In monkeys, the absorption of [^{14}C]pentachlorophenol after a single oral dose of 10 mg/kg was evaluated by measuring the plasma concentration of pentachlorophenol (Braun and Sauerhoff, 1976). Peak plasma concentrations (10-30 μg/g) were attained within 12-24 hours after oral administration, but there were sex differences in the rate constants and half-lives of absorption. In males, the absorption rate constant is 0.215 hours^{-1}, with a half-life of 3.64 hours; in females, the absorption rate constant is 0.383 hours^{-1}, with a half-life of 1.81 hours.

Oral administration of a single dose of 0.1 mg/kg pentachlorophenol to four male volunteers resulted in an absorption half-life of 1.3±0.4 hours (Braun et al., 1978), which followed first-order kinetics.

Distribution. Ninety-six hours after intraperitoneal injection of ~15-37 mg/kg of [^{14}C]pentachlorophenol to female NMRI mice, the highest concentrations (weight of pentachlorophenol/organ weight) of pentachlorophenol were found in the gall bladder and liver (Jakobson and Yllner, 1971). As indicated in Table 6, smaller concentrations were identified in the stomach and intestines. The brain, heart, lung and kidney—the only additional organs examined—had low concentrations of pentachlorophenol. One week after dosing, the concentration of pentachlorophenol remained elevated in the liver, but the concentration was not reported for the gall bladder or gastrointestinal tract.

After oral administration of a single dose of 10 or 100 mg/kg of [^{14}C]pentachlorophenol to male and female Sprague-Dawley rats, the liver and kidneys contained the greatest concentrations of radioactivity; the brain, spleen and fat contained the lowest concentrations (Braun et al., 1977). The other organs analyzed included lungs, testes, ovaries, heart, skeletal fat, perirenal fat and adrenal glands. The peak plasma concentrations of pentachlorophenol occurred 4–6 hours after dosing. The concentration of radioactivity was usually higher in the organs and plasma of the females than it was in the organs and plasma of the males.

Two female rhesus monkeys were killed and the radioactivity was measured in the major organs 360 days after the oral administration of a single dose of 10 mg/kg [^{14}C]pentachlorophenol (Braun and Sauerhoff, 1976). The liver, the small intestines and the large intestines contained the largest fraction of the administered radioactivity, 1.41±0.54%, 2.60±1.86% and 5.00±2.91%, respectively.

Metabolism. Analysis of the urine intraperitoneal injection of ~15-37 mg/kg of [^{14}C]pentachlorophenol to female NMRI mice indicated that ~33% of the administered radioactivity was present as unmetabolized pentachlorophenol. In addition to free pentachlorophenol, conjugated pentachlorophenol and tetrachlorohydroquinone were identified in the urine of the treated mice (Jakobson and Yllner, 1971).

After a single oral dose of 100 mg/kg of [^{14}C]pentachlorophenol to Sprague-Dawley rats, analysis of the urine indicated that 48% of the administered radioactivity was present as unmetabolized pentachlorophenol and 10% as tetrachlorohydroquinone. The remaining urinary radioactivity, 6% of the administered dose, appeared to be conjugated pentachlorophenol (Braun et al., 1977).

Analysis of the urine of NMRI mice and Sprague-Dawley rats after oral administration of 25 mg/kg of [$_{14}$C]pentachlorophenol indicated that 41–43% of the urinary radioactivity in both species was present as unchanged pentachlorophenol and that 5 and 24% of the urinary radioactivity in the rats and mice, respectively, was tetrachlorohydroquinone (Ahlborg et al., 1974). Ahlborg et al. (1974) also reported that tetrachlorohydroquinone was identified in the urine of humans occupationally exposed to pentachlorophenol.

Braun and Saurhoff (1976) analyzed the urine from monkeys given a single oral dose of 10 mg/kg of [^{14}C]pentachlorophenol and found that essentially all the radioactivity was present as pentachlorophenol. Analysis of the urine from four human males after ingestion of a single oral dose of 0.1 mg/kg of pentachlorophenol indicated that both unmetabolized and conjugated pentachlorophenol were present (Braun et al., 1978).

Excretion. After intraperitoneal injection of ~15-37 mg/kg of [^{14}C]pentachlorophenol to female NMRI mice, ~50% of the administered radioactivity was excreted in the urine within the first 24 hours (Jakobson and Yllner, 1971). As the administered dose increased, the percentage of the administered radioactivity excreted in the urine within 24 hours of dosing increased (Table 7). The amount of radioactivity excreted in the feces ranged from 3.8-11.5% of the administered [^{14}C]pentachlorophenol 96-168 hours after dosing. Of the administered radioactivity, <0.05% was detected in the expired air.

After administration of a single oral dose of 10 or 100 mg/kg of [^{14}C]pentachlorophenol to three male and three female Sprague-Dawley rats, urine and feces were collected at 24-hour intervals, and the rats killed after 8 or 9 days (Braun et al., 1977). Collection of the expired air at 12-hour intervals from the rats treated with 10 mg/kg of [^{14}C]pentachlorophenol indicated that <1% of the administered dose was excreted as CO_2; the expired air from the high-dose group rats was not analyzed. Of the administered radioactivity, ~80 and 64%, respectively, were recovered from the urine of rats within 8 or 9 days after receiving single oral doses of 10 or 100 mg/kg. Approximately 19 and 34% of the administered radioactivity were recovered, respectively, from the feces of rats within 8 or 9 days after receiving single oral doses of 10 or 100 mg/kg. Elimination of pentachlorophenol in the urine and feces was biphasic in the male and female rats receiving 10 mg/kg [^{14}C]pentachlorophenol, with half-lives of 17 and 13 hours, respectively, in the α phase and 13 and 27 hours, respectively, in the β phase. At the high-dose level, elimination was biphasic for the males only with a half-life of 40 hours in the α phase and 33 hours in the β phase. In the high-dose females, elimination followed first-order kinetics with a half-life of 27 hours. Both the binding of pentachlorophenol to plasma proteins and the retention of pentachlorophenol in the liver appeared to be responsible for the β phase of elimination.

Ahlborg et al. (1974) intraperitoneally injected NMRI mice and Sprague-Dawley rats with 25 mg/kg of [^{14}C]pentachlorophenol; 96 hours after administration, ~70% of the administered radioactivity was recovered from the urine of both rats and mice. Within 96 hours after oral administration of the same dose, however, only 45 and 25% of the administered radioactivity were recovered from the urine of rats and mice, respectively.

TABLE 7
Excretion of the Activity After Intraperitoneal Injection of ^{14}C-Pentachlorophenol in Female NMRI Mice*

Dose (mg/kg)	Percentage of Dose in Urine					Percentage of Dose in Feces
	0-24 hours	24-48 hours	48-72 hours	72-96 hours	96-168 hours	
14.8	44.6	30.0	5.0	3.4	NR	0.1 (0-24 hours) 7.7 (24-96 hours)
18.2	45.4	15.9	6.1	4.6	NR	4.2 (0-24 hours) 0.8 (24-96 hours)
37.2	54.3	12.3	4.2	2.3	NR	3.8 (0-96 hours)
35.2	59.7	11.1	5.8	3.1	1.8	6.5 (0-24 hours) 3.2 (24-168 hours)
36.8	58.9	14.9	4.3	1.4	0.9	1.5 (0-168 hours)

*Source: Jakobson and Yllner, 1971
NR = Not reported

After oral administration of 31–40 mg/kg of [^{14}C]pentachlorophenol to four male and four female rats, ~65% of the administered radioactivity was recovered in the urine within 72 hours of dosing (Larsen et al., 1972). Larsen et al. (1972) suggested that urinary excretion was biphasic in rats, with half-lives of 10 hours for the α phase and 102 days for the β phase. Fecal elimination of radioactivity was not measured and recovery of the administered dose from the urine was assumed to approach 100%.

After the administration of a single oral dose of 10 mg/kg [^{14}C]pentachlorophenol to three male and three female rhesus monkeys, the plasma, urine and feces were collected over a period ranging from 132-360 hours (Braun and Sauerhoff, 1976). Approximately 10-20% of the administered radioactivity was excreted in the feces at a slow but steady rate, indicating that enterohepatic circulation was occurring. Approximately 70-80% of the administered radioactivity was recovered from the urine. Urinary excretion followed first-order kinetics with a half-life of ~92.4 hours in females and 40.8 hours in males.

Approximately 74 and 12% of the administered dose were identified in the urine as pentachlorophenol glucuronide, respectively, 168 hours after administration of a single oral dose of 0.1 mg/kg of pentachlorophenol to four male volunteers (Braun et al., 1978). Braun et al. (1978) described the kinetics of human elimination of pentachlorophenol as first-order in a one-compartment open-system model with substantial enterohepatic circulation. The half-life for urinary elimination of pentachlorophenol is 33.1 ± 5.4 hours, and for urinary elimination of pentachlorophenol glucuronide 12.7 ± 5.4 hours.

Casarett et al. (1969) studied the urinary excretion of pentachlorophenol by workers at a wood treatment plant in Hawaii and found that there may be a difference in excretion kinetics between acute high-level and chronic low-level exposures. In two workers exposed by inhalation to pentachlorophenol for 45 minutes, urinary excretion half-lives were ~10 hours. Following chronic exposures, however, concentrations in the urine were decreased by only ~60–80% in workers absent from work for ≤18 days. The authors suggested that this difference may be due to a "four-compartment system for chronic inhalation exposure consisting of lung, plasma, proteins, urine and tissue depots."

Summary. Pentachlorophenol is readily absorbed from the gastrointestinal tract of rats, mice, monkeys and humans (Braun et al., 1977, 1978; Ahlborg et al., 1974; Braun and Sauerhoff, 1976). Peak plasma concentrations are reached within 12–24 hours after oral administration to monkeys (Braun and Sauerhoff, 1976), but 4–6 hours after oral administration to rats (Braun et al., 1977). After oral administration, the highest concentration of radioactivity was found in the liver and gastrointestinal tract of monkeys (Braun and Sauerhoff, 1976), and in the liver and kidneys of rats (Braun et al., 1977). In rats and mice, tetrachlorohydroquinone was identified in the urine (Jakobson and Yllner, 1971; Braun et al., 1977; Ahlborg et al., 1974), as well as unmetabolized pentachlorophenol and glucuronide-conjugated pentachlorophenol. Al-

though Ahlborg et al. (1974) reported that oxidative dechlorination of pentachlorophenol occurs in humans, as evidenced by the presence of tetrachlorohydroquinone in the urine of workers occupationally exposed (probably by inhalation), analysis of human urine after ingestion of pentachlorophenol revealed the presence of conjugated pentachlorophenol and unmetabolized pentachlorophenol (Braun et al., 1978).

The primary route of excretion after oral administration in all species studied is in the urine (Braun et al., 1977, 1978; Ahlborg et al., 1974; Larsen et al., 1972; Braun and Sauerhoff, 1976). Although urinary excretion followed second-order kinetics in rats (Larsen et al., 1972; Braun et al., 1977) except in females receiving a single high dose (100 mg/kg) of pentachlorophenol, urinary excretion of pentachlorophenol in humans and monkeys followed first-order kinetics (Braun and Sauerhoff, 1976; Braun et al., 1978). Enterohepatic circulation played an importation role in the pharmacokinetics of pentachlorophenol. The half-life of pentachlorophenol in the plasma is longer in female rats and monkeys than it is in male rats and monkeys (Braun et al., 1978; Braun and Sauerhoff, 1976).

EFFECTS

Carcinogenicity. The carcinogenicity of pentachlorophenol has been evaluated after oral exposure (BRL, 1968; Innes et al., 1969; Schwetz et al., 1977), subcutaneous injection (BRL, 1968; Catalina et al., 1981) and skin-painting experiments (Boutwell and Bosch, 1959). An increased incidence of cancer was not reported in any of these studies. Only the oral studies, however, will be summarized in detail because the other routes of administration are not relevant to this study.

Groups of 18 male and 18 female mice from each strain, (C57BL/6 x C3H/Anf)F1 and C57BL/6 x AKR)F1, were treated by gavage with 46.4 mg/kg/day of pentachlorophenol (Dowicide 7®) on days 7-28 of age. Subsequently, the mice were exposed to 130 ppm of dietary Dowicide 7® for the next 78 weeks. Groups of mice from each strain that received either no treatment or the vehicle only (gelatin) served as controls. Although not reported in the carcinogenicity study (Innes et al., 1969; BRL, 1968) itself, Schwetz et al. (1977) reported that the commercial Dowicide EC-7® contained 90% pentachlorophenol, 10% tetrachlorophenol, 22.55 ppm dibenzo-p-dioxins, 6.2 ppm dibenzofurans and 400 ppm hexachlorobenzene. The parameters evaluated included survival rate and tumor incidence. There was no significant difference between the incidence of tumors in the pentachlorophenol treated group and the control group.

In the 22- to 24-month study by Schwetz et al. (1977), Sprague-Dawley rats exposed to diets providing 1, 3, 10 or 30 mg/kg/day of commercial pentachlorophenol consisting of 90% pentachlorophenol, 10% tetrachlorophenol, 22.44 ppm dibenzo-p-dioxin, 6.2 ppm dibenzofuran and 400 ppm hexachlorobenzene had no significant difference in tumor incidence compared with control rats.

TABLE 8
Mutagenicity of Pentachlorophenol in Microbial Systems*

Assay	Species/Strain	Application	Dose	Purity	Result	S-9	Comments	Reference
Reverse mutation	S. typhimurium 8 NS	spot	NA	commercial	-	-	NC	Anderson et al., 1972
Reverse mutation	S. typhimurium TA1535, TA1537, TA1538, TA98 and TA100	plate incorporation	up to 5 mg/plate	NS	-/-	+/-	NC	Simmon et al., 1977
Reverse mutation	S. typhimurium TA1535, TA1537, TA1538, TA98 and TA100	liquid suspension	NS	NS	-/-	+/-	NC	Simmon et al., 1977
Reverse mutation	S. typhimurium TA1535, TA1537, TA1538, TA98 and TA100	vapor	NA	NS	-/-	+/-	NC	Simmon et al., 1977
Reverse mutation	S. typhimurium TA100, TA98, TA1535, TA1537 and TA1538	plate incorporation	up to 5 mg/plate	commercial	-/-	+/-	NC	Moriya et al., 1983
Reverse mutation	S. typhimurium TA1535, TA1537, TA1538, TA98 and TA100	plate incorporation	up to 10 mg/plate	technical	-/-	+/-	NC	Waters et al., 1982
Reverse mutation	S. marcescens a21, a742	spot	NA	NS	-	-	NC	Fahrig, 1974
Reverse mutation	E. coli WP2uvr-	plate incorporation	up to 5 mg/plate	NS	-/-	+/-	NC	Simmon et al., 1977
Forward mutation	E. coli NS	liquid suspension	NS	NS	-	-	NC	Fahrig, 1974
Reverse mutation	E. coli WP2hcr	plate incorporation	up to 5 mg/plate	NS	-/-	+/-	NC	Moriya et al., 1983
Reverse mutation	E. coli WP2uvrA	plate incorporation	up to 10 mg/plate	technical	-/-	+/-	NC	Waters et al., 1982

TABLE 8 (continued)

Assay	Species/Strain	Application	Dose	Purity	Result	S-9	Comments	Reference
Rec assay	E. coli p3478, W3100	spot	up to 10 mg	technical	-	-	NC	Waters et al., 1982
Rec assay	B. subtillis M45, H17	spot	up to 10 mg/plate	technical	+	-	NC	Waters et al., 1982
Rec assay	B. subtillis H17Rec+, M45rec-	spot	5 µg/plate	NS	+	-	weakly positive results	Shirasu, 1976
Mitotic recombination	S. cerevisiae D3	liquid suspension	NS	NS	-/-	+/-	NC	Simmon et al., 1977
Mitotic recombination or forward mutation	S. cerevisiae MP-1	liquid suspension	400 mg/l	purified	+	-	positive for forward mutation, positive for intragenic and negative for intergenic recombination	Fahrig et al., 1977
Mitotic gene conversion	S. cerevisiae ade2, trp5	liquid suspension	0.19 mM	NS	+	-	NC	Fahrig, 1974

*Source: U.S. EPA, 1984a

NA = Not applicable; NS = not specified; NC = no comment

According to U.S. EPA (1981b), the use of pentachlorophenol poses a carcinogenic risk because it is contaminated with hexachlorodibenzo-p-dioxin and hexachlorobenzene. Technical grade pentachlorophenol and Dowicide EC-7 are being tested by the NTP in long-term feeding studies in mice (NTP, 1985). The chronic histopathology of the study of Dowicide EC-7 is in progress. The quality assessment of the study on technical pentachlorophenol is in progress.

Mutagenicity. As indicated in Table 8, neither commercial nor technical pentachlorophenol is mutagenic to *Salmonella typhimurium* (Anderson et al., 1972; Simmon et al., 1977; Lemma and Ames, 1975; Moriya et al., 1983; Waters et al., 1982; Buselmaier et al., 1973) or *Escherichia coli* (Simmon et al., 1977; Fahrig, 1974; Moriya et al., 1983; Waters et al., 1982) with or without metabolic activation. There was no increase in sex-linked recessive lethal mutation in *Drosophila melanogaster* treated with pentachlorophenol (Vogel and Chandler, 1974). Technical pentachlorophenol was mutagenic to *Bacillus subtilis* at a concentration of 10 mg/plate (Waters et al., 1982). Earlier, Shirasu (1976) reported weakly positive results when *B. subtilis* was treated with pentachlorophenol of unspecified purity at concentrations up to 5 μg/plate. Fahrig et al. (1977) tested pentachlorophenol *in vitro* with the multipurpose strain MP-1 of *Saccharomyces cerevisiae* for simultaneous detection of forward mutations, intergenic or intragenic recombinations. Pentachlorophenol at a concentration of 400 mg/ℓ was active in the mutation system as well as intragenic recombination system. While tested in an *in vivo* mammalian spot test system, pentachlorophenol at a concentration of 50 mg/kg showed positive response. Comparing the frequency of spots induced with the dose applied, pentachlorophenol exhibited weak mutagenic activity, while activity of pentachloro predioxin, an impurity associated with the manufacturing process, can be compared with that of dimethylnitrosamine (DMN). This indicates that pentachlorophenol has definite but weak mutagenic activity while pentachloro predioxin has much stronger mutagenic activity.

Intraperitoneal injection of female C57BL/6 JHan mice with 50 or 100 mg/kg purified pentachlorophenol on day 10 of gestation resulted in a slightly increased incidence of mutation in the pups, as evidenced by coat-color spots (Fahrig et al., 1977). Although not statistically significant, Wyllie et al. (1975) reported an increased number of breaks and gaps in the chromosomes of cultivated peripheral blood cells from workers exposed to an average monthly air concentration of pentachlorophenol that ranged from 263.0-1887.9 mg/m^3.

Teratogenicity. There were six studies located in the available literature that provided data on the teratogenicity and fetotoxicity of pentachlorophenol, four gavage studies (Courtney et al., 1976; Schwetz et al., 1974; Hinkle, 1973; Larsen et al., 1975; Larsen, 1976) and two dietary studies (Exon and Koller, 1982; Schwetz et al., 1977). Because many preparations of pentachlorophenol are contaminated with highly toxic substances, such as dioxins, special attention must be paid to the purity of the pentachlorophenol tested. In studies where technical and purified pentachlorophenol have both been evaluated (Schwetz et al., 1974, 1977), only the results of the experiments using purified pentachlorophenol will be reported in detail.

In the Courtney et al. (1976) study, a group of seven pregnant C-D rats was treated by gavage on days 7-18 of gestation with 75 mg/kg/day of pentachlorophenol in corn oil:acetone (9:1). The purity of the pentachlorophenol was not reported. The control group of five pregnant C-D rats was treated by gavage on days 7-18 of gestation with the vehicle of corn oil:acetone (9:1). The experiment was terminated 1 or 2 days before parturition. The parameters evaluated included maternal and fetal body weights, maternal liver-to-body weight ratio, litter size (average number of live fetuses/litter), fetal mortality rate (number of dead or resorbed fetuses/total number of fetuses) and teratogenicity (number of abnormal fetuses/litter). Decreased fetal body weight was the only statistically significant effect on the endpoints evaluated. Histological evaluation of the fetuses was not reported.

In the Schwetz et al. (1974) study, groups of 20-40 Sprague-Dawley rats were treated by gavage with 0, 5, 15, 30 or 50 mg/kg/day of pentachlorophenol in corn oil on days 6-15 of gestation. The study was terminated on day 21 of gestation. The purified pentachlorophenol contained >98% pentachlorophenol, 0.32% other chlorophenols, 0.5% higher chlorinated phenoxyphenols, <2ppm dibenzo-p-dioxins and <1.5 ppm dibenzofurans. In rats given purified pentachlorophenol, the parameters evaluated included maternal body weight, fetal body weight and crown-rump length, fetal resorptions, fetal sex ratio and incidence of fetal anomalies (subcutaneous edema, dilated ureter, and skeletal variations). There was a 100% resorption rate (229/229 fetuses; 19/19 litters) in the rats treated with 50 mg/kg/day of pentachlorophenol, and a 97.5% resorption rate (233/239 fetuses; 20/30 litters) in the rats treated with 30 mg/kg/day. Maternal body weights were significantly depressed compared with controls on day 21 of gestation in the groups receiving doses of 30 or 50 mg/kg/day. Fetal body weights were significantly depressed at 30 mg/kg/day. Compared with control values, the fetal crown-rump length was also significantly decreased in the group receiving 30 mg/kg/day. At 30 mg/kg/day pentachlorophenol, 83% of the pups were males, indicating that the female pups were relatively more sensitive to the pentachlorophenol effects. Resorption rates and male:female ratios were not affected in the rats treated with 5 or 15 mg/kg/day. Dilated ureters were not observed in any of the fetuses, but the incidence of subcutaneous edema was significantly increased in the fetuses of dams treated with 15 or 30 mg/kg/day pentachlorophenol. The incidence of all types of fetal skeletal anomalies, including delayed ossification of the skull, lumbar spurs, supernumerary (lumbar or fused) ribs, vertebral and sternebral anomalies (supernumerary, abnormal shape, missing, delayed or unfused centers of ossification), was significantly increased at the 15 mg/kg/day dose level. Although only 3/5 types of skeletal anomalies (ribs, vertebrae and sternebrae) were observed at the 30 mg/kg/day dose level, only 2/20 litters tested at this dose level produced viable offspring. At 5 mg/kg/day, the lowest dose tested, cranial ossificaton was delayed significantly, but no other skeletal abnormalities were reported.

A group of CD rats (number not specified) was treated by gavage with a single dose of 60 mg/kg pentachlorophenol (purity and concentration of contaminants not specified)

on day 8, 9, 10, 11 or 13 of gestation (Larsen, 1976; Larsen et al., 1975). The dams were sacrificed on day 20. No significant effect on fetal resorptions was observed; however, fetal weight was significantly reduced in the offspring of dams treated on day 9 or 10. There were four malformed fetuses in the treated groups: a dwarf in the group treated on day 8, and one tailless, one macrophthalmia and one exencephaly in the group treated on day 9. No malformed fetuses were found in the control group. Because the number of fetuses examined was not reported, it is not known if the malformations are statistically or toxicologically significant.

As reviewed by U.S. EPA (1978) and U.S. EPA (1982), Hinkle (1973) administered pentachlorophenol (purity and concentration of contaminants not specified) by gavage to Golden Syrian hamsters at doses of 0, 1.25, 2.5, 5, 10 or 20 mg/kg/day on days 6-10 of gestation. No differences between control and experimental groups were found for maternal body weight, fetal weight, litter size or number of resorptions. The investigator stated that the feed of the control hamsters was contaminated with detectable (not otherwise specified) amounts of pentachlorophenol.

After 62 days of ingesting diet providing 0, 3 or 30 mg/kg/day of a test substance containing pentachlorophenol, groups of 10 male and 20 female Sprague-Dawley rats were mated (Schwetz et al., 1977). The test substance (Dowicide EC-7) consisted of 90% pentachlorophenol, 10% tetrachlorophenol, 22.0 ppm dibenzo-p-dioxin, 6.0 ppm dibenzofuran and 400.0 ppm hexachlorobenzene. The rats were maintained on their respective diets during mating, gestation and lactation. The dams and pups were killed 21 days after birth of the pups, and the males were killed after 169 days of treatment. The parameters evaluated included body weights of adults and pups, fertility rate (number of pregnant females/number of mated females), number of pups born alive, pup survival rate (number of pups alive 1, 7, 14 and 21 days postpartum/number of pups born), and the number of litters with skeletal variations in weanling rats. At 30 mg/kg/day, there were significant decreases in the body weights of dams 21 days postpartum and in the pups from birth through weaning, in the number of live born pups, and in the survival rate of pups 7, 14 and 21 days after birth but not during the first 24 hours after birth. The fertility rate was not affected at 30 mg/kg/day, but the number of skeletal variations, such as lumbar spurs or vertebrae with unfused centers, was significantly increased. Schwetz et al. (1977) concluded that the increased incidence of skeletal variations in the high-dose group was indicative of fetotoxicity rather than teratogenicity. The decreased survival of the pups may reflect an adverse reproductive effect. No effects on the parameters evaluated were observed in rats ingesting 3 mg/kg/day pentachlorophenol.

Groups of 12-20 female Sprague-Dawley rats were exposed to 0, 5, 50 or 500 ppm dietary pentachlorophenol beginning at 21 days of age (Exon and Koller, 1982). The rats were mated at 90 days of age and were maintained on their pentachlorophenol diets through mating, gestation and lactation. The experiment was terminated when the pups were 21 days old. Analysis of the pentachlorophenol indicated that the test chemical consisted of 85.5% pentachlorophenol, 7.4% tetrachlorophenols and 408 ppm

dioxins. The parameters evaluated included fertility rate, litter size, birthweight, number of stillborn pups/total number of pups, weaning weight and hematology of the pups at weaning (red and white cell counts, hemoglobin levels, packed cell volume and mean corpuscular volume). Although the litter size was reduced in the rats exposed to 500 ppm dietary pentachlorophenol, the decrease was not statistically significant. No other effects were reported.

Other Reproductive Effects. Pertinent data regarding other reproductive effects of pentachlorophenol could not be located in the available literature as cited in the Appendix.

Chronic and Subchronic Toxicity. There are six studies available that provide data pertinent for quantitative risk assessment of the chronic and subchronic toxicity of pentachlorophenol. Because many preparations of pentachlorophenol are contaminated with small but measurable amounts of highly toxic substances, such as dioxins, special attention must be paid to the purity of the pentachlorophenol tested. In studies where technical and purified pentachlorophenol have been evaluated (Goldstein et al., 1977; Kimbrough and Linder, 1978; Knudsen et al., 1974; Johnson et al., 1973; Kerkvliet et al., 1982), only the results of the experiments using purified pentachlorophenol will be reported in detail.

Groups of 27 male and 27 female Sprague-Dawley rats were exposed to pentachlorophenol at dietary levels that provided doses of 0, 1, 3, 10 or 30 mg/kg/day (Schwetz et al., 1977). The test chemical (Dowicide EC-7) consisted of 90% pentachlorophenol, 10% tetrachlorophenol, 22.55 ppm dibenzo-p-dioxin, 6.2 ppm dibenzofuran and 400.0 ppm hexachlorobenzene. The female rats were maintained on pentachlorophenol-contaminated diets for 24 months, but because of decreased survival rate in the experimental and control groups, the male rats were sacrificed after 22 months of treatment. The parameters evaluated included body weight, food consumption, hematology (12 months and terminal), urinalysis (12 months and terminal), clinical chemistry, organ weights (brain, heart, liver, kidneys and testes), and gross and microscopic evaluation of unspecified organs. Food consumption, hematology and organ weights were unaffected by treatment. Mortality rate data were not given. At 30 mg/kg/day, the body weight gain of female rats was significantly reduced and SGPT levels were significantly increased in both males and females. Gross observations revealed darkly discolored livers and kidneys in ~50% of the females ingesting 30 mg/kg/day pentachlorophenol; microscopic evaluation revealed brown pigment in the hepatocytes of 16/27 females but only 1/27 males, and in the renal tubular epithelial cells of 19/27 females. At 10 mg/kg/day, darkly discolored livers (2/27) and kidneys (3/27) were grossly observed in the female rats. Brown pigment was identified microscopically in hepatocytes of 8/27 females and in the renal tubular epithelial cells of 7/27 females. No other compound-related effects were reported in the rats ingesting 10 mg/kg/day of pentachlorophenol. No effects were reported at 3 mg/kg/day.

Groups of 10 male and 10 female Sherman rats were fed diets containing 0, 20, 100 or 500 ppm purified pentachlorophenol for 8 months (Kimbrough and Linder, 1978). The purified pentachlorophenol consisted of >99% pentachlorophenol and <0.1 ppm dibenzofurans or dibenzodioxins. The parameters evaluated included body and organ weights, food consumption and microscopic examination of brain, lung, heart, spleen, liver, kidneys, thyroid, parathyroid, gastrointestinal tract, adrenal, bladder and reproductive organs. The body weights of rats exposed to 500 ppm purified pentachlorophenol in the diet were decreased, the weight of males being affected more than females. A significant compound-related, but not dose-related, increase in kidney weights of all the male rats exposed to purified pentachlorophenol was observed, but there were no remarkable histopathological renal lesions. Histological changes, including enlarged centrilobular hepatocytes in eight males and five females and pigmented Kupffer cells in three females, were reported in the livers of rats exposed to 500 ppm purified pentachlorophenol in the diet. No other lesions were reported. Although hepatic porphyria in female rats and extensive hepatic histopathology, including fibrosis, vacuolization and hyperplasia were observed in both male and female rats exposed to technical pentachlorophenol, these changes were not observed in the rats exposed to purified pentachlorophenol. The technical pentachlorophenol was contaminated with unspecified amounts of dibenzodioxins and dibenzofurans.

Groups of six female Sherman rats were exposed to dietary concentrations of 0, 20, 100 or 500 ppm pure or technical pentachlorophenol for 8 months (Goldstein et al., 1977). The pure preparation consisted of >99% pentachlorophenol; no dibenzo-p-dioxins or dibenzofurans (detection limit 0.1 ppm) were identified in this preparation. The technical preparation consisted of 84.6% pentachlorophenol and 3% tetrachlorophenol, and was contaminated with 1908 ppm dibenzo-p-dioxins and 790 ppm dibenzofurans. The parameters evaluated included body weight, food consumption, liver weight, urinary porphyrins and precursors and hepatic enzyme levels (δ-aminolevulinic acid synthetase, aryl hydrocarbon hydroxylase, aminopurine N-demethylase, p-nitrophenol glucuronyl transferase, cytochrome P-450 and total microsomal heme). The body weight of the rats exposed to 500 ppm dietary pure pentachlorophenol was depressed. No treatment-associated effect was observed on food consumption, liver weight or liver-to-body weight ratio at any of the dose levels. The urinary porphyrin and porphyrin precursor levels were not affected by pure pentachlorophenol, although a significant increase was observed in the urine of rats exposed to 100 or 500 ppm of technical pentachlorophenol. The only hepatic enzyme level affected by pure pentachlorophenol was p-nitrophenol glucuronyl transferase, which increased in rats exposed at the 500 ppm level. Significant increases in the levels of all the hepatic enzymes tested, except aminopurine N-demethylase, were observed in the rats exposed to technical pentachlorophenol.

Groups of 10 male and 10 female weanling rats (strain unspecified) were exposed for 12 weeks to 0, 25, 50 or 200 ppm dietary pentachlorophenol contaminated with 200 ppm of octachlorodibenzo-p-dioxin and 82 ppm of pre-octachloro-dibenzo-p-dioxin (Knudsen et al., 1974). The parameters evaluated included food consumption, body

TABLE 9
Oral LD_{50} Values for Pentachlorophenol

Species	Sex	LD_{50} (mg/kg)	Purity	Reference
Rat (adult)	NS	150	commercial (~90%)	Schwetz et al., 1977
Rat (3- to 4-day-old)	NS	65	commercial (~90%)	Schwetz et al., 1977
Rat	NS	27-100	commercial	Knudsen et al., 1974
Rat	NS	300	90%	Nishimura et al., 1980
Rat	M	156-269	commercial (~88%)	Schwetz et al., 1974
Rat	F	97-188	commercial (~88%)	Schwetz et al., 1974
Mice	NS	120-140	commercial	Knudsen et al., 1974
Mice	M	125-252	>99%	Borzelleca, 1983; Borzelleca et al., 1984
Mice	F	65-212	>99%	Borzelleca, 1983; Borzelleca et al., 1984
Rabbit	NS	100-130	commercial	Knudsen et al., 1974
Dog	NS	150-200	commercial	Knudsen et al., 1974
Guinea pig	NS	~100	commercial	Knudsen et al., 1974

NS = Not specified

and organ weights, hematology, clinical chemistry (SGPT, but not SGOT), urinalysis, liver enzyme levels and histopathology of the major organs in the control and high-dose groups. Only the liver and kidneys were evaluated microscopically in the low- and middle-dose groups. Although the decreased body weight of the female rats was statistically significant in only the low- and the high-dose groups, the biological significance of the body weight changes in the female rats is unclear. Similarly, the biological significance of statistically significant increases in the serum alkaline phosphatase levels in female rats exposed to 25 and 200, but not 50 ppm and the significantly increased

serum urea levels in females exposed to 25, but not 50 or 200 ppm is not apparent. Dose-related significantly increased relative liver weight was observed in females at 50 ($p<0.01$) and the 200 ppm ($p<0.001$) compared with controls. The level of aryl hydrolase was significantly increased in the high-dose females, but the incidence of centrilobular vacuolization was not different from controls. In the male rats, hepatic centrilobular vacuolization (5/10) and increased microsomal enzyme activity occurred at 200 ppm, and hepatic centrilobular vacuolization (4/10) occurred at 50 ppm. The male rats exposed to 50 or 200 ppm had significantly decreased hemoglobin levels and red blood cell counts at the end of the experiment; the hemoglobin levels and red blood cell counts were significantly increased in these dose groups at the mid point of the experiment. Serum glucose levels were significantly increased in the male rats in the high-dose group.

In a study designed to test the toxicity of two preparations (commercial and purified) of pentachlorophenol, Johnson et al. (1973) exposed groups of Sprague-Dawley rats (number/sex/group unspecified) to diets providing doses of 0, 3, 10 or 300 mg/kg/day for 90 days. The commercial pentachlorophenol preparations consisted of only 85-90% pentachlorophenol and contained up to 1% caustic insolubles and 584-2537 ppm dioxins, and the "purified" pentachlorophenol preparation contained 88-93% pentachlorophenol and up to 31 ppm of dioxins. The content of caustic insolubles was not reported. The parameters evaluated included body weight, food consumption, hematology, clinical chemistry, urinalysis, terminal organ weight and histology of unspecified tissues. Kidney weights were significantly increased in the rats ingesting 30 mg/kg/day, and liver weight was significantly increased in the rats ingesting 10 to 30 mg/kg/day purified pentachlorophenol. Histopathological lesions were not observed in the kidneys or livers of the rats exposed to any level of purified pentachlorophenol, and no effects were reported at 3 mg/kg/day, the same no effect level reported in the Schwetz et al. (1977) 2-year dietary study. Exposure to all three dose levels of the commercial pentachlorophenol preparation, however, resulted in increased liver kidney weights, which was associated with microscopic hepatic pathology at the highest dose tested.

Several immunological tests were performed on male C57BL/6(B6) mice after exposure to 0, 50 or 500 ppm of dietary pure (>99%) or technical grade (86%) pentachlorophenol for 10-12 weeks (Kerkvliet et al., 1982). The purified pentachlorophenol had no adverse effect on immunosurveillance as measured by the ability of the male mice to resist challenge with a 3-methylcholanthrene-induced syngeneic transplantable tumor. In tests of immunocompetence using MSV-induced tumors, groups of male mice (~10-16/group) were inoculated with MSV, which produced transient subcutaneous tumors at the site of injection over a period of 15 weeks. The mice were then challenged with an injection of MSV-transformed cancer cells. Although the mice ingesting purified pentachlorophenol did not develop progressive secondary tumor growth (i.e., the injected cells were prevented from forming tumors in the pentachlorophenol-treated hosts), histological examination of the spleens of the mice ingesting purified pentachlorophenol 7 weeks after injection of cancer cells indicated that 2/9 mice in the

low-dose group and 4/9 mice in the high-dose group had splenic cancer. None of the 13 mice in the control group had cancer. The authors interpreted their results as indicating that ingestion of purified pentachlorophenol decreased the immunocompetence of male mice. Application of the Fisher Exact test to the data demonstrated that the decrease was statistically significant only at the high-dose level. Exposure of mice to both dietary levels of the technical pentachlorophenol, however, had a profound enhancing effect on tumor susceptibility.

Other Relevant Information. Several oral LD_{50} values are listed in Table 9. The LD_{50} value for females is usually lower than for males.

Summary. Because many preparations of pentachlorophenol are contaminated with small but measurable amounts of highly toxic substances, such as dibenzodioxins, special attention must be paid to the composition of the pentachlorophenol solution tested. In studies where technical and purified pentachlorophenol have both been evaluated (Schwetz et al., 1974; Goldstein et al., 1977; Kimbrough and Linder, 1978; Knudsen et al., 1974; Johnson et al., 1973; Kerkvliet et al., 1982), only the results of the experiments using purified pentachlorophenol were reported in detail. Oral exposure to pentachlorophenol was not carcinogenic in mice (BRL, 1968; Innes et al., 1969) or rats (Schwetz et al., 1977), regardless of the composition of the pentachlorophenol solution tested. Although there are a few studies that suggest pentachlorophenol may be mutagenic in *B. subtilis* (Waters et al., 1982; Shirasu, 1976), in yeast, *Saccharomyces cerevisiae* (Fahrig et al., 1977) and in mice, as evidenced by the coat-color spot test (Fahrig et al., 1977), no evidence of mutagenicity was reported in *S. typhimurium* (Anderson et al., 1972; Simmon et al., 1977; Lemma and Ames, 1975; Moriya et al., 1983; Waters et al., 1982; Buselmaier et al., 1973) or in *E. coli* (Simmon et al., 1977; Fahrig, 1974; Moriya et al., 1983; Waters et al., 1982) with or without metabnolic activation.

Three teratogenicity/reproductive toxicity studies (Schwetz et al., 1974, 1977; Courtney et al., 1976) indicate that pentachlorophenol is fetotoxic in rats at oral dose levels $\geq$5 mg/kg/day. At the highest dose tested (500 ppm) in a fourth teratogenicity/reproductive toxicity study (Exon and Koller, 1982), there was a decrease in litter size, which was not statistically significant. The lowest dose tested (5 mg/kg/day) by Schwetz et al. (1977) was the lowest dose at which any evidence of fetotoxicity, as indicated by delayed ossification, was observed. No adverse fetal or reproductive effects were reported at $\leq$ 3 mg/kg/day (Schwetz et al., 1977; Exon and Koller, 1982). In subchronic and chronic toxicity studies, adverse effects occurred primarily in the liver (Kerkvliet et al., 1982; Johnson et al., 1973; Knudsen et al., 1974; Goldstein et al., 1977; Kimbrough and Linder, 1978; Schwetz et al., 1977), the kidney (Johnson et al., 1973; Kimbrough and Linder, 1978; Schwetz et al., 1977) and the immune system (Kerkvliet et al., 1982). Knudsen et al. (1974) reported increased liver weights in female rats and centrilobular vacuolization in male rats exposed to diets containing $\geq$ 50 ppm commercial pentachlorophenol, which contained 282 ppm dioxins. In the remaining

TABLE 10
Acute Toxicity of Pentachlorophenol to Freshwater Fishes and Amphibians

Species	Concentration (µg/ℓ)	Effect	Reference
Rainbow trout *Salmo gairdneri*	157	48-hour LC_{50}	Alabaster, 1957
	52	96-hour LC_{50}	Johnson and Finley, 1980
	200	48-hour LC_{50}	Sloof et al., 1983
	170	NOLC	Sloof et al., 1983
	70	detection limit, change in respiratory frequency	Sloof, 1979
	66	96-hour LC_{50}	Dominguez and Chapman, 1984
	46	41-day LC_{100}	Chapman, 1969
Brown trout *Salmo trutta*	197	24-hour LC_{50}	Ribo and Kaiser, 1983
	157	48-hour LC_{50}	Alabaster, 1957
Atlantic salmon *Salmo salar*	46	24-hour, altered temperature preference	Peterson, 1976
Chinook salmon *Oncorhynchus tschawytscha*	68	96-hour LC_{50}	Johnson and Finley, 1980
Coho salmon *Oncorhynchus kisutch*	65	96-hour LC_{50}, NaPCP	Spehar et al., 1981
Brook trout *Salvelinus fontinalis*	109	336-hour LC_{50}	Cardwell et al., 1976
Carp *Cyprinus carpio*	9.5	96-hour LC_{50}, larvae	Verma et al., 1981
	0.5-0.6	estimated MATC	Verma et al., 1981

TABLE 10 (continued)

Species	Concentration (µg/ℓ)	Effect	Reference
Medaka	1100	48-hour LC_{50}	Sloof et al., 1983
Oryzias latipes	930	NOLC	Sloof et al., 1983
Goldfish	270	25-hour LC_{50}	Kobayashi et al., 1979
Carassius auratus	117-191	96-hour LC_{50}	Phipps and Holcombe, 1985
Zebrafish	400	LC_{50}	Sloof, 1979
Brachydanio rerio			
Bluegill	115-152	96-hour LC_{50}	Phipps and Holcombe, 1985
Lepomis macrochirus	32	96-hour LC_{50}	Johnson and Finley, 1980
	300	96-hour LC_{50}	Pierce, 1978
	174	336-hour LC_{50}	Cardwell et al., 1976
Largemouth bass	136-287	96-hour LC_{50}	Johansen et al., 1985
Micropterus salmoides	50	7 days, decreased food consumption and feeding efficiency	Mathers et al., 1985
Channel catfish	54	96-hour LC_{50}	Phipps and Holcombe, 1985
Ictalurus punctatus	68	96-hour LC_{50}	Johnson and Finley, 1980
Fathead minnow	141	336-hour LC_{50}	Cardwell et al., 1976
Pimphales promelas	220-230	96-hour LC_{50}	Phipps et al., 1981
	200-220	192-hour LC_{50}	Phipps et al., 1981
	210	48-hour LC_{50}	Sloof et al., 1983
	170	NOLC	Sloof et al., 1983
	205	96-hour LC_{50}	Johnson and Finley, 1980
	470	96-hour LC_{50}	Cleveland et al., 1982
	223-286	96-hour LC_{50}	Phipps and Holcombe, 1985

TABLE 10 (continued)

Species	Concentration (µg/ℓ)	Effect	Reference
Guppy	45-160	96-hour LC_{50}	Adema and Vink, 1981
Poecilia reticulata	36-110	7-day LC_{50}	Adema and Vink, 1981
	850	48-hour LC_{50}	Sloof et al., 1983
	620	NOLC	Sloof et al., 1983
	372	7- to 14-day LC_{50}	Ribo and Kaiser, 1983
	970	96-hour LC_{50}, PCP	Gupta et al., 1982a
	770	96-hour LC_{50}, NaPCP	Gupta et al., 1982a
	200	24-hour LC_{50}	Tanaka et al., 1981
	45-1150	96-hour LC_{50}	Adema and Vink, 1981
Catfish	290	96-hour LC_{50}	Verma et al., 1980
Saccobranchus fossilis	16	safe concentration	Verma et al., 1980
Gouramy	450	96-hour LC_{50}	Verma et al., 1980
Colisa fasciatus	5.1	safe concentration	Verma et al., 1980
Knifefish	32-131	96-hour LC_{50}	Gupta et al., 1982b
Notopterus notopterus	1.1-4.5	safe concentrations	Gupta et al., 1982b
	83	96-hour LC_{50}	Verma et al., 1980
	22	safe concentrations	Verma et al., 1980
Amphibians	260	48-hour LC_{50}	Sloof et al., 1983
Xenopus laevis	210	NOLC	Sloof et al., 1983
Ambystoma mexicanus	300	48-hour LC_{50}	Sloof et al., 1983
	130	NOLC	Sloof et al., 1983

studies, increased liver weight (Johnson et al., 1973) and increased pigmentation of hepatocytes (Schwetz et al., 1977) were observed at oral doses of ≥10 mg/kg/day (~90%), and SGPT levels significantly increased in rats ingesting 30 mg/kg/day pentachlorophenol (~90%) for 2 years (Schwetz et al., 1977). Increased kidney weight unaccompanied by renal histopathology was reported in rats exposed to a dietary concentration ≥20 ppm of pentachlorophenol (>99%) for 8 months (Kimbrough and Linder, 1978) and in rats ingesting 30 mg/kg/day (~90%) for 90 days (Johnson et al., 1973). Increased pigmentation of the renal tubular epithelial cells was reported in rats ingesting 10 or 30 mg/kg/day pentachlorophenol for 2 years (Schwetz et al., 1977). Although decreased immunocompetence was reported in mice exposed to dietary levels of 50 or 500 ppm of pentachlorophenol (>99%) for 34 weeks (Kerkvliet et al., 1982), the decrease was statistically significant only at the higher dose.

AQUATIC TOXICITY

Acute. U.S. EPA (1980a) presented a large volume of information concerning effects of pentachlorophenol on aquatic organisms. Pentachlorophenol was acutely toxic to freshwater fish species, with 96-hour LC_{50} values ranging from 34-600 μg/ℓ . Salmonids were generally more sensitive than other species with LC_{50} values of 34-128 μg/ℓ compared with LC_{50} values of 60-600 μg/ℓ for nonsalmonid species. The most sensitive salmonid species was coho salmon, *Oncorhynchus kisutch,* with an LC_{50} value of 34 μg/ (U.S. EPA, 1980a). The most sensitive nonsalmonid fish species was the bluegill, *Lepomis macrochirus,* with an LC_{50} value of 60-77 μg/ℓ.

Data from studies not reported in U.S. EPA (1980a) are presented in Table 10, and these support the generalization that salmonids are more sensitive than other freshwater species. Some nonsalmonid freshwater species also have been found to have LC_{50} values <100 μg/ . Among these are carp, *Cyprinus carpio,* larvae with an LC_{50} of 9.5 μg/ (Verma et al., 1981), the bluegill, *L. macrochirus,* with an LC_{50} of 32 μg/ (Johnson and Finley, 1980), the channel catfish, *Ictalurus punctatus,* with an LC_{50} of 54-68 μg/ (Johnson and Finley, 1980; Phipps and Holcombe, 1985) and the knifefish, in *Notopterus notopterus,* with an LC_{50} of 83 μg/ (Verma et al., 1980).

Limited information was available for freshwater invertebrates (U.S. EPA, 1980a). EC_{50} values were 240-800 μg/ℓ for *Daphnia magna* and 2000 μg/ℓ for *Daphnia pulex.* The worm, *Tubifex tubifex,* had 24-hour LC_{50} values of 286-1294 μg/ℓ. U.S. EPA (1980a) concluded that these limited data indicated that freshwater invertebrates were about as sensitive to pentachlorophenol as nonsalmonid fish species. Data from more recent studies support this generalization (Table 11).

Among the saltwater fish species discussed by U.S. EPA (1980a), the most sensitive was the pinfish, *Lagodon rhomboides,* prolarvae, with an LC_{50} value of 38 μg/ℓ (Borthwick and Schimmel, 1978). No data from more recent studies (Table 12) indicated a more sensitive species, although the goby, *Gobius minutus,* had a 48-hour and 7-day

TABLE 11
Acute Toxicity of Pentachlorophenol to Freshwater Fishes and Marine Invertebrates

Species	Concentration (µg/ℓ)	Effect	Reference
		CRUSTACEANS	
Daphnia sp.	1,492	24-hour LC_{50}	Ribo and Kaiser, 1983
Daphnia magna	170-510	21-day LC_{50}	Adema, 1978
	1,500	24-hour LC_{50}	Leblanc, 1980
	680	96-hour LC_{50}	Leblanc, 1980
	320	NOEC	Leblanc, 1980
	480	48-hour LC_{50}	Sloof et al., 1983
	160	NOLC	Sloof et al., 1983
	330	EC_0, immobilization	Bringmann and Kuehn, 1982
	800	EC_{50}, immobilization	Bringmann and Kuehn, 1982
	1,800	EC_{100}, immobilization	Bringmann and Kuehn, 1982
	500-510	24-hour EC_{50}	Berglind and Dave, 1984
	370-440	48-hour EC_{50}	Berglind and Dave, 1984
	143	48-hour LC_{50}	Mount and Norberg, 1984
	1,050	48-hour LC_{50}, larvae	Adema and Vink, 1981
	1,400	48-hour LC_{50}, adult	Adema and Vink, 1981
	800	96-hour LC_{50}, larvae	Adema and Vink, 1981
	680	96-hour LC_{50}, adult	Adema and Vink, 1981
	600	7-day LC_{50}, larvae	Adema and Vink, 1981
	400	14- to 21-day LC_{50}, larvae	Adema and Vink, 1981
	580	7-day LC_{50}, adult	Adema and Vink, 1981
Daphnia pulex	2,000	48-hour LC_{50}	Sloof et al., 1983
	1,200	NOLC	Sloof et al., 1983
	246	48-hour LC_{50}	Mount and Norberg, 1984
Daphnia cucullata	1,500	48-hour LC_{50}	Sloof et al., 1983

TABLE 11 (continued)

Species	Concentration (µg/ℓ)	Effect	Reference
		CRUSTACEANS (cont.)	
Ceriodaphnia reticulata	164	48-hour LC_{50}	Mount and Norberg, 1984
Simocephalus vetulus	217	48-hour LC_{50}	Mount and Norberg, 1984
Chaetogammarus marinus	550	96-hour LC_{50}, larvae	Adema and Vink, 1981
	450	96-hour LC_{50}, adult	Adema and Vink, 1981
	210	14-day LC_{50}, larvae and adult	Adema and Vink, 1981
	180	21-day LC_{50}, larvae and adult	Adema and Vink, 1981
Temora longicornis	200	48-hour LC_{50}	Adema and Vink, 1981
	170	96-hour LC_{50}	Adema and Vink, 1981
Decapod			
Crangon crangon	1,790	96-hour LC_{50}, adults (NaPCP)	VanDijk et al., 1977
	112	96-hour LC_{50}, larvae (NaPCP)	VanDijk et al., 1977
	10,000	96-hour LC_{50}	Adema and Vink, 1981
	5,000	7-day LC_{50}	Adema and Vink, 1981
Palaemon elegans	10,390	96-hour LC_{50}, adults (NaPCP)	VanDijk et al., 1977
	84	96-hour LC_{50}, larvae (NaPCP)	VanDijk et al., 1977
Palaemonetes varians	5,090	96-hour LC_{50}, adults (NaPCP)	VanDijk et al., 1977
	363	96-hour LC_{50}, larvae (NaPCP)	VanDijk et al., 1977
	7,500	96-hour LC_{50}	Adema and Vink, 1981
	5,800	7-day LC_{50}	Adema and Vink, 1981
Amphipod			
Gammarus pulex	700	48-hour LC_{50}	Sloof, 1983

TABLE 11 (continued)

Species	Concentration (µg/ℓ)	Effect	Reference
		CRUSTACEANS (cont.)	
Isopod Asellus aquaticus	2,900	48-hour LC_{50}	Sloof, 1983
Copepod Pseudodiaptomus coronatus	68	96-hour LC_{50} (NaPCP)	Hauch et al., 1980
Brine shrimp Artemia salina	4,600	96-hour LC_{50}, larvae	Adema and Vink, 1981
	16,000	96-hour LC_{50}, adult	Adema and Vink, 1981
	4,500	7-day LC_{50}, larvae	Adema and Vink, 1981
	11,000	7-day LC_{50}, adult	Adema and Vink, 1981
Grass shrimp Palaemonetes pugio	515	96 hours, no significant mortality	Schimmel et al., 1978
	473	9 days, 50% reduction in limb regeneration	Rao et al., 1978
Brown shrimp Penaeus aztecus	195	96 hours, no significant mortality	Schimmel et al., 1978
		INSECTS	
Chironomid Chironomus gr. thummi	110	48-hour LC_{50}	Sloof, 1983
Corixid Corixa punctata	1,100	48-hour LC_{50}	Sloof, 1983

TABLE 11 (continued)

Species	Concentration (µg/ℓ)	Effect	Reference
		INSECTS (cont.)	
Odonate Ischnura elegans	4,200	48-hour LC_{50}	Sloof, 1983
Stonefly Nemoura cinerca	380	48-hour LC_{50}	Sloof, 1983
Mayfly Cloeon dipterum	5,900	48-hour LC_{50}	Sloof, 1983
Mosquito Aedes aegypti	7,200 1,800	48-hour LC_{50} 48-hour NOLC	Slooff et al., 1983 Slooff et al., 1983
Mosquito Culex pipiens	34,000 24,000	48-hour LC_{50} 48-hour NOLC	Slooff et al., 1983 Slooff et al., 1983
		MOLLUSKS	
Snail Lymnaea acuminata	160 190	96-hour LC_{50} 96-hour LC_{50} (NaPCP)	Gupta and Rao, 1982 Gupta and Rao, 1982
Snail Lymnaea stagnalis	560 200 240 180	48-hour LC_{50} 48-hour NOLC 96-hour LC_{50} 16-day LC_{50}	Slooff et al., 1983 Slooff et al., 1983 Adema and Vink, 1981 Adema and Vink, 1981
Snail Lymnaea luteola	112 575	96-hour LC_{50} 96-hour LC_{50} (NaPCP)	Gupta et al., 1984 Gupta et al., 1984

TABLE 11 (continued)

Species	Concentration (µg/ℓ)	Effect	Reference
		OTHERS	
Polychaete *Neanthes virens*	365,720	acutely lethal	Carr and Neff, 1981
Ophyrotrocha diadema	620	96-hour LC_{50}, larvae	Adema and Vink, 1981
	1,200	96-hour LC_{50}, adult	Adema and Vink, 1981
	280	28- to 41-day LC_{50}, larvae	Adema and Vink, 1981
	300	28- to 30-day LC_{50}, adult	Adema and Vink, 1981
Oligochaete *Limnodrilus hoffmeisteri*	330	96-hour LC_{50} (NaPCP)	Chapman et al., 1982
Oligochaete *Tubifex tubifex*	380	96-hour LC_{50} (NaPCP)	Chapman et al., 1982
Tubificidae	1,000	48-hour LC_{50}	Slooff, 1983
Leech *Erpobdella octoculata*	250	48-hour LC_{50}	Slooff, 1983
Leech	3,500	24-hour LC_{50}	Kulkarni et al., 1984
Poechilobdella viridis	1000-3000	effect on oxygen consumption	Kulkarni et al., 1984
Flatworm *Pugesia lugubris*	130	48-hour LC_{50}	Slooff, 1983
Hydra	730	48-hour LC_{50}	Slooff et al., 1983
Hydra oligactis	430	48-hour NOLC	Slooff et al., 1983
Ciliate protozoan *Colpidium campidum*	600	minimum active dose	Dive et al., 1980

Species	Concentration (µg/ℓ)	Effect	Reference
		MOLLUSKS (cont.)	
Snail Viviparus bengalensis	840 720	96-hour LC_{50} 96-hour LC_{50} (NaPCP)	Gupta and Durve, 1984 Gupta and Durve, 1984
Snail Onchidium verruculatum	355	96-hour LC_{50}	Deshpande and Nagabhushanam, 1981
Mussel Mytilus edulis	18,000 950 400 750	96-hour LC_{50} 7-day LC_{50} 48-hours, 22-69% abnormal larvae 14-day LC_{50}	Adema and Vink, 1981 Adema and Vink, 1981 Dimick and Breese, 1965 Dimick and Breese, 1965
Mussel Dreissana polymorpha	110-190 52-180 42-140	96-hour LC_{50} 7-day LC_{50} 14-day LC_{50}	Dimick and Breese, 1965 Dimick and Breese, 1965 Dimick and Breese, 1965
Gastropod Crepidula fornicata	1,200 460	48-hour LC_{50} 7-day LC_{50}	Dimick and Breese, 1965 Dimick and Breese, 1965
Pacific oyster Crassostrea gigas	55	48-hours, 62% abnormal embryos	Woelke, 1972
Eastern oyster Crassostrea virginica	250	48 hours, no embryos developed	Davis and Hidu, 1962
Carpet shell Tapes phillippinarum	100	120 hours, lethal	Tomiyama et al., 1962

TABLE 12
Acute Toxicity of Pentachlorophenol to Marine Fishes

Species	Concentration (µg/ℓ)	Effect	Reference
Mullet Mugil cephalus	100	48-hours 33% mortality	Thomas et al., 1981
	200	24-hours 33% mortality	Thomas et al., 1981
Flounder Pleuronectes platessa	60-140	96-hour LC_{50}, larvae	Adema and Vink, 1981
	140-170	96-hour LC_{50}, juveniles	Adema and Vink, 1981
	750	96-hour LC_{50}, eggs	Adema and Vink, 1981
	100-130	7-day LC_{50}, juveniles	Adema and Vink, 1981
	50	8-week LC_{50}, eggs	Adema and Vink, 1981
Goby Gobius minutus	45	48-hour and 7-day LC_{50}	Adema and Vink, 1981
Longnose killifish Fundulus similis	306	96-hours, no significant mortality	Schimmel et al., 1978

LC_{50} value of 38 µg/ℓ, and eggs and larvae of the flounder, *Pleuronectes platessa*, had LC_{50} values of 50 and 60 µg/ℓ, respectively (Adema and Vink, 1981).

Of the marine invertebrates discussed by U.S. EPA (1980a), the most sensitive was the Eastern oyster, *Crassostrea virginica*, with an LC_{50} value of 40 µg/ℓ (Borthwick and Sehimmel, 1978). None of the more recent studies cited in Table 11 identify a more sensitive species; however, Van Dijk et al. (1977) reported that larval stages of crustaceans are 10-100 times more sensitive than adults. From the data presented in Table 12 and U.S. EPA (1980a), the most sensitive (i.e., $LC_{50} < 200$ µg/ℓ) invertebrate species are the Eastern oyster, the larvae of the crustaceans, *Crangon crangon* and *Palaemon elegans* (VanDijk et al., 1977), and the copepod, *Pseudodiaptomus coronatus* (Hauch et al., 1980) among the marine species, and the chironomid, *Chironomus gr. thummi* (Slooff, 1983), and the snail, *Lymnaea luteola* (Gupta et al., 1984), among the freshwater species.

Pentachlorophenol was more acutely toxic to aquatic organisms at lower pH, probably because low pH favors the formation of molecular pentachlorophenol, which is more toxic than the ionized form that is favored at higher pH (U.S. EPA, 1980a). Other possible explanations are that pentachlorophenol degradation is accelerated at higher pH (Dalela et al., 1980) or that pentachlorophenol uptake is slower at higher pH (Kobayashi and Kishino, 1980). Data from studies not cited in U.S. EPA (1980a) confirm the trend of increasing pentachlorophenol toxicity at lower pH (Table 13).

Chronic. The available information concerning chronic toxicity of pentachlorophenol is summarized in Tables 14 and 15. In fathead minnow embryo-larval tests, growth was a more sensitive response to pentachlorophenol than survival and was inhibited at 45 μg/ℓ in one study (Holcombe et al., 1982) and 13 μg/ℓ in another (Cleveland et al., 1982). Growth was also more sensitive than survival in guppies, with a NOEC for growth inhibition of <180 μg/ℓ (Adema and Vink, 1981), and in salmonids, with growth inhibition in several species occurring at 3.2-28 μg/ℓ (U.S. EPA, 1980a). In the nonsalmonid fish species tested, mortality was generally a more sensitive indicator of chronic pentachlorophenol toxicity than growth or reproductive parameters (see Table 14). The lowest reported threshold for chronic toxicity in freshwater fish species was 1.8 μg/ℓ (NaPCP), which inhibited food conversion efficiency and growth of sockeye salmon, *Oncorhynchus nerka* (Webb and Brett, 1973). The lowest reported threshold in a saltwater fish species was 10 μg/ℓ, which was a NOEC for growth inhibition, mortality and malformations in the flounder, *Pleuronectes platessa* (Adema and Vink, 1981). Pentachlorophenol caused increased mortality of the sheepshead minnow, *Cyprinodon variegatus*, at 88 μg/ , but growth and fecundity were unaffected at concentrations up to 195 μg/ℓ (Parrish et al., 1978).

The most sensitive freshwater invertebrate species was the hydra, *Hydra littoralis*, with a threshold for inhibition of multiplication of 19 μg/ℓ (Dannenberg, 1984). In a 21-day test, chronic mortality of *Daphnia magna* occurred at 320 μg/ but not 180 μg/ℓ, while reproduction was unaffected at these levels (Adema, 1978). Among the marine invertebrates tested, the Eastern oyster, *Crassostrea virginica*, was quite sensitive, with larval survival and growth affected at 50 μg/ℓ (Davis and Hidu, 1969). The polychaete worm, *Ophryotrocha diadema*, was also quite sensitive; its reproduction was inhibited at 23-75 μg/ℓ (Adema and Vink, 1981).

As was the case with acute toxicity, chronic toxicity of pentachlorophenol was affected by pH. As measured in a fathead minnow embryo-larval test, toxicity increased with decreasing pH. Survival (pH 6.5, 8.5) and growth (pH 7.5, 8.0) were the most sensitive effects dependent on pH. Chronic values based on these effects were as follows: pH 6.5, 24 μg/ℓ ; pH 7.5, 4.0 μg/ℓ ; pH 8.0, 49 μg/ℓ ; and pH 8.5, 89 μg/ℓ (Spehar et al., 1985).

Plants. The available information concerning the effects of pentachlorophenol on aquatic plants is summarized in Table 16. It is difficult to draw conclusions from these data because of the large variability in reported toxic levels. For example, there is a 1000-fold difference in toxic concentrations reported for *Chlorella pyrenoidona;* Adema and Vink (1981) reported a 96-hour EC_{50} for growth inhibition of 7000 μg/ℓ, while

TABLE 13
Effect of pH on Pentachlorophenol Toxicity

Species	pH	Concentration (µg/l)	Effect	Reference
Goldfish	5.5	52	24-hour LC_{50}	Kobayashi and Kishino, 1980
Carassius auratus	6.0	60	24-hour LC_{50}	Kobayashi and Kishino, 1980
	7.0	82	24-hour LC_{50}	Kobayashi and Kishino, 1980
	8.0	250	24-hour LC_{50}	Kobayashi and Kishino, 1980
	9.0	2,200	24-hour LC_{50}	Kobayashi and Kishino, 1980
	10.0	16,000	24-hour LC_{50}	Kobayashi and Kishino, 1980
Fathead minnow	6.5	95	96-hour LC_{50}	Spehar et al., 1985
Pimephales promelas	7.5	218	96-hour LC_{50}	Spehar et al., 1985
	8.0	261	96-hour LC_{50}	Spehar et al., 1985
	8.5	378	96-hour LC_{50}	Spehar et al., 1985
Colisa fasciatus	4.6	92	96-hour LC_{50}	Dalela et al., 1980
	6.0	340	96-hour LC_{50}	Dalela et al., 1980
	7.3	450	96-hour LC_{50}	Dalela et al., 1980
	8.8	850	96-hour LC_{50}	Dalela et al., 1980
Saccobranchus fossilis	4.6	62	96-hour LC_{50}	Dalela et al., 1980
	6.0	180	96-hour LC_{50}	Dalela et al., 1980
	7.3	290	96-hour LC_{50}	Dalela et al., 1980
	8.8	750	96-hour LC_{50}	Dalela et al., 1980
Notopterus notopterus	4.6	4.3	96-hour LC_{50}	Dalela et al., 1980
	6.0	50	96-hour LC_{50}	Dalela et al., 1980
	7.3	83	96-hour LC_{50}	Dalela et al., 1980
	8.8	370	96-hour LC_{50}	Dalela et al., 1980

TABLE 13 (continued)

Species	pH	Concentration (µg/ℓ)	Effect	Reference
Guppy	5.0	42	96-hour LC_{50}	Saarikoski and Viluksela, 1981
Poecilia reticulata	6.0	117	96-hour LC_{50}	Saarikoski and Viluksela, 1981
	6.1	128	96-hour LC_{50}	Konnemann and Musch, 1981
	7.0	442	96-hour LC_{50}	Saarikoski and Viluksela, 1981
	7.3	376	96-hour LC_{50}	Konnemann and Musch, 1981
	7.8	767	96-hour LC_{50}	Konnemann and Musch, 1981
	8.0	911	96-hour LC_{50}	Saarikoski and Viluksela, 1981
Carp	5.5	28	24-hour LC_{50}	Khangarot et al., 1985
Cyprinus carpio	6.5	44	24-hour LC_{50}	Khangarot et al., 1985
	7.5	208	24-hour LC_{50}	Khangarot et al., 1985
	8.5	400	24-hour LC_{50}	Khangarot et al., 1985
	9.5	3,126	24-hour LC_{50}	Khangarot et al., 1985
Zebrafish	<6	10-20	48-hour LC_{50}	Dave, 1984
Brachydanio rerio	7-8	100-200	48-hour LC_{50}	Dave, 1984
Amphipod	6.5	92	96-hour LC_{50}	Spehar et al., 1985
Gammarus pseudolimnaeus	7.5	121	96-hour LC_{50}	Spehar et al., 1985
	8.0	484	96-hour LC_{50}	Spehar et al., 1985
	8.5	790	96-hour LC_{50}	Spehar et al., 1985
Amphipod	6.5	139	96-hour LC_{50}	Spehar et al., 1985
Crangonyx pseudogracilis	7.5	465	96-hour LC_{50}	Spehar et al., 1985
	8.0	929	96-hour LC_{50}	Spehar et al., 1985
	8.5	1,344	96-hour LC_{50}	Spehar et al., 1985
Worm	7.5	286	24-hour LC_{50}	Whitley, 1968
Tubifex tubifex	8.5	619	24-hour LC_{50}	Whitley, 1968
	9.5	1,294	24-hour LC_{50}	Whitley, 1968

TABLE 14
Chronic and Subchronic Toxicity of Pentachlorophenol to Fishes

Species	Concentration (µg/L)	Duration (days)	Effect	Reference
			FRESHWATER	
Fathead minnow	128	28	decreased survival, embryo-larval test	Holcombe et al., 1982
Pimephales promelas	73	28	decreased growth, embryo-larval test	Holcombe et al., 1982
	45	28	NOEC, embryo-larval test	Holcombe et al., 1982
	200-330	NR	threshold for long-term mortality	Alexander et al., 1983
	24-89	32	chronic values, pH 6.5-8.5	Spehar et al., 1985
	13	90	reduced growth, threshold, commercial PCP	Cleveland et al., 1982
	27	90	reduced survival, threshold, commercial PCP	Cleveland et al., 1982
	85	90	reduced growth, threshold, purified PCP	Cleveland et al., 1982
Guppy	580	NR	threshold for long-term mortality	Alexander et al., 1983
Poecilia reticulata	<180	60	NOEC for growth inhibition	Adema and Vink, 1981
Rainbow trout	10-20	NR	threshold for decreased biomass production from newly fertilized eggs exposed through hatching and 4 weeks after yolk sac absorption	Hodson and Blunt, 1981
Salmo gairdneri	60	72	interocular hemorrhage, fry	Dominguez and Chapman, 1984
	34	72	pre-hatch mortality significantly increased	Dominguez and Chapman, 1984
	19	72	alevin mortality 3 times that of controls	Dominguez and Chapman, 1984
	11	72	no adverse effects	Dominguez and Chapman, 1984
	14	72	chronic toxicity threshold	Dominguez and Chapman, 1984
	7.4-28	20-92	9-27% growth inhibition	Chapman, 1969; Chapman and Shumway, 1978; Matida et al., 1970
Sockeye salmon	3.2	NR	10% growth inhibition	Webb and Brett, 1973
Oncorhynchus nerka	1.8	14-60	threshold for effects on growth, food conversion efficiency, NaPCP	Webb and Brett, 1973
Salmonids	3.2-28.0	21-91	10-27% growth inhibition	U.S. EPA, 1980a
Bluegill	100	32	histological effects	Owen and Rosso, 1981
Lepomis macrochirus				
			SALTWATER	
Sheepshead minnow	88	151	increased mortality	Parrish et al., 1978
Cyprinodon variegatus	195	151	reduced hatching success	Parrish et al., 1978
	47	151	no adverse effects	Parrish et al., 1978
Flounder	10	56	NOEC for growth inhibition, mortality, malformations	Adema and Vink, 1981
Pleuronectes platessa	50	56	LC_{50}	Adema and Vink, 1981

NR = Not reported

TABLE 15
Chronic and Subchronic Toxicity of Pentachlorophenol to Invertebrates

Species	Concentration (µg/L)	Duration (days)	Effect	Reference
			FRESHWATER	
Waterflea				
Daphnia magna	340	21	NOEC for reproductive effects	Adema and Vink, 1981
	320	21	mortality	Adema, 1978
	180	21	no mortality	Adema, 1978
Snail				
Lymnaea stagnalis	130	16	EC_{50} (morphological defects and hatching)	Adema and Vink, 1981
	50	16	NOEC	Adema and Vink, 1981
Rotifer				
Brachionus rubens	50-200	NR	effects on populations	Halbach et al., 1983
Hydra				
Hydra littoralis	19	NR	threshold for inhibition of multiplication	Dannenberg, 1984
			SALTWATER	
Oyster				
Crassostrea virginica	100	14	0% survival of larvae	Davis and Hidu, 1969
	50	14	decreased larval survival and growth	Davis and Hidu, 1969
	25	14	no effect	Davis and Hidu, 1969
	34	8	EC_{50}, reduced shell deposition	Schimmel et al., 1978
Brine shrimp				
Artemia salina	5800	28	NOEC for reproductive effects	Adema and Vink, 1981
	4400	28	LC_{50}	Adema and Vink, 1981
Crustacean				
Chaetogammarus marinus	100	56	NOEC for growth inhibition	Adema and Vink, 1981
	180	56	LC_{50}	Adema and Vink, 1981
Polychaete worm				
Ophryotrocha diadema	3	NR	NOEC for inhibition of reproduction	Hooftmann and Vink, 1980
	23-75	41	EC_{50} for inhibition of reproduction	Adema and Vink, 1981
	10-13	41	NOEC for inhibition of reproduction	Adema and Vink, 1981
Neanthes virens	100	60	decreased glycogen reserves, increased ascorbic acid content of parapodial tissue	Carr and Neff, 1981

NR = Not reported

TABLE 16
Effects of Pentachlorophenol on Aquatic Plants

Species	Concentration (µg/ℓ)	Effect	Reference
Chlorella pyrenoidosa	7000	96-hour EC_{50}, growth inhibition	Adema and Vink, 1981
	7.5	72-hours, complete destruction of chlorophyll	Huang and Gloyna, 1968
Chlorella ovalis	5500	96-hour EC_{50}, growth inhibition	Adema and Vink, 1981
Scenedesmus quadricauda	80	96-hour EC_{50}, growth inhibition	Adema and Vink, 1981
Scenedesmus abliquus	133-266	toxicity threshold	Senger and Ruhl, 1980
Selenastrum capricornutum	53-133	toxicity threshold	Jayaweera et al., 1982
	410	48-hour EC_{50}, growth inhibition	Crossland and Wolf, 1985
	290	96-hour EC_{50}, growth inhibition	Crossland and Wolf, 1985
Chlamydomonas reinhardi	266	3-minute EC_{50}, inhibition of oxidative phosphorylation	Crossland and Wolf, 1985
Chlamydomonas sp.	1400	96-hour EC_{50}, growth inhibition	Adema and Vink, 1981
Lemna minor	190	48-hour EC_{50}, chlorosis	Blackman et al., 1955
	800	48-hour EC_{50}, chlorosis	Blackman et al., 1955
	200	inhibits photosynthetic oxygen production	Huber et al., 1982
Macrocystis pyrifera	1000	inhibits photosynthesis	Crossland and Wolf, 1985
Monochrysis lutheri	293	12 days, 58% decrease in cell numbers	Woelke, 1965
Monochrysis sp.	200	96-hour EC_{50}, growth inhibition	Adema and Vink, 1981
Skeletonema cosfatum	17-20	EC_{50}, decrease in cell numbers	U.S. EPA, 1980a
Thalassiosira pseudonana	179-205	EC_{50}, decrease in cell numbers	U.S. EPA, 1980a
Dunaliella tertiolecta	170-206	EC_{50}, decrease in cell numbers	U.S. EPA, 1980a
Dunaliella sp.	3600	96-hour EC_{50}, growth inhibition	Adema and Vink, 1981
Thalassia testudinum	740	EC_{50}, inhibition of oxygen production	Walsh et al., 1982
Phaeodactylum tricornutum	3000	96-hour EC_{50}, growth inhibition	Adema and Vink, 1981
Enteromorpha sp.	50-100	30 days, lethal, experimental ponds	Crossland and Wolf, 1985
Chara sp.	50-100	30 days, lethal, experimental ponds	Crossland and Wolf, 1985
Algae	2000	21 days, 50% growth inhibition	McKee and Wolf, 1963
	125-2000	inhibition of photosynthesis, estuarine phytoplanteton	Erickson and Hawkins, 1980

Huang and Gloyna (1968) reported that 7.5 μg/ℓ caused death in 72 hours. The data presented in Table 16 do not provide evidence that aquatic plants are more tolerant or sensitive to pentachlorophenol than fish or invertebrates. Therefore, concentrations protective of one of these groups should be protective of plants as well (U.S. EPA, 1980a).

Residues. Results of pentachlorophenol uptake and depuration studies with aquatic organisms are summarized in Table 17. These studies also indicate that peak pentachlorophenol accumulation or steady-state generally occurs in 4-8 days' exposure in many species including killifish (Trujillo et al., 1982), bluegills (Pruitt et al., 1977), goldfish (Kobayashi and Akitake, 1975) and oysters (Schimmel et al., 1978). Trujillo et al. (1982) suggested that this amount of time was required for the induction of enzymes to detoxify and excrete pentachlorophenol. The data of Pruitt et al. (1977) also indicate that some detoxification mechanism is induced because tissue pentachlorophenol concentrations were lower after 16 days' exposure than 8 days' exposure, even though ambient pentachlorophenol concentrations were constant.

Several investigators have studied elimination of pentachlorophenol by various species. Kobayashi and Nakamura (1979) found that in goldfish exposed to 100 μg/ℓ for 24 hours, 52% of the absorbed pentachlorophenol was excreted through the gills, 30% of which was free pentachlorophenol. The rest of the absorbed pentachlorophenol was excreted by the kidney (24%) and bile (22%), primarily in the form of pentachlorophenol-glucuronide and pentachlorophenol-sulfate. Glickman et al. (1977) reported that bile of exposed rainbow trout contained high concentrations of pentachlorophenol, mostly as glucuronide conjugates. Lu and Metcalf (1975) concluded that pentachlorophenol conjugation was a significant detoxification mechanism in *Daphnia,* mosquitos (*Culex*) and snails (*Physa*), but not in mosquitofish (*Gambusia affinis*), which stored unconjugated pentachlorophenol. Kobayashi et al. (1969) reported that pentachlorophenol in shellfish was stored as the parent compound, either free or conjugated.

The data presented in Table 17 indicate that pentachlorophenol is not strongly bioaccumulated by aquatic organisms, with steady-state BCF values generally <1000. Of all the species tested, the marine polychaete worm, *Lanice conchilega,* accumulated pentachlorophenol to the greatest extent, with a BCF of 3830 (Ernst, 1979).

Studies show that there is a relationship between pentachlorophenol uptake and toxicity. Kobayashi and Kishino (1980) observed that tissues of goldfish poisoned by pentachlorophenol contained 82-115 μg/g (average 92 μg/g) and suggested that fish would die when the body burden of pentachlorophenol reached 92 μg/g, regardless of the pentachlorophenol concentration in the water. Spehar et al. (1985) reported similar results for fathead minnows exposed to pentachlorophenol; dead fish contained pentachlorophenol at an average concentration of 75 μg/g, while the highest concentration observed in unaffected fish was 35 μg/g. The results of these two studies indicate that there is a threshold tissue pentachlorophenol level that must be reached for toxic effects to occur.

TABLE 17
Results of Pentachlorophenol Uptake and Depuration Studies with Aquatic Organisms

Species	Water Concentration (µg/L)	Exposure Duration	Tissue	Tissue Concentration (ppb)	BCF	Elimination $t_{1/2}$	Reference
			FRESHWATER				
Goldfish	200	NR	whole fish	95	475	NR	Kobayashi et al., 1979
Carassius auratus	200	NR	whole fish	116,000	580	NR	Kobayashi and Akitake, 1975
	100	120 hours	whole fish	100,000	1000	10 hours	Kobayashi and Akitake, 1975
Rainbow trout	25	24 hours	liver	16,000	640	9.8 hours	Glickman et al., 1977
Salmo gairdneri	25	24 hours	blood	6500	260	6.2 hours	Glickman et al., 1977
	25	24 hours	fat	6000	240	23 hours	Glickman et al., 1977
	25	24 hours	muscle	1000	40	6.9 hours	Glickman et al., 1977
	0.66	115 days	organs	2200	3333	7 days	Niimi and McFadden, 1982
	0.66	115 days	whole fish	160	242	7 days	Niimi and McFadden, 1982
	0.035	115 days	whole fish	10-20	286-572	7 days	Niimi and McFadden, 1982
	10 or 40 ppb diet	110 days	whole fish	1-2	NR	7 days	Niimi and Cho, 1983
	3 ppm diet	110 days	whole fish	20-40	NR	7 days	Niimi and Cho, 1983
Bluegill	100	8 days	muscle	1300	13	NR	Pruitt et al., 1977
Lepomis macrochirus	100	8 days	gill	6000	60	NR	Pruitt et al., 1977
	100	8 days	digestive tract	21,000	210	NR	Pruitt et al., 1977
	100	8 days	liver	35,000	350	NR	Pruitt et al., 1977
Fathead minnow	11.1	32 days	whole fish	NR	770	NR	Veith et al., 1979b
Pimephales promelas	3.1-451	32 days	whole fish	2500-69,000	281-1066	NR	Spehar et al., 1985
Carp	10-50	72 hours	blood	4000	NR	NR	Gluth et al., 1985
Cyprinus carpio	10-50	72 hours	liver, kidney, intestine, gills	2000-3000	NR	NR	Gluth et al., 1985
	10-50	72 hours	muscle	<500	NR	NR	Gluth et al., 1985
Mosquitofish	8	NR	NR	1080	132	NR	Lu and Metcalf, 1975; Lu et al., 1978
Gambusia affinis							
Snail	8	NR	NR	1140	21	NR	Lu and Metcalf, 1975; Lu et al., 1978
Physa sp.							
Mosquito	8	NR	NR	170	26	NR	Lu and Metcalf, 1975; Lu et al., 1978
Culex sp.							
Waterflea	8	NR	NR	1170	205	NR	Lu and Metcalf, 1975; Lu et al., 1978
Daphnia sp.							
Igae	8	NR	NR	11	5	NR	Lu and Metcalf, 1975; Lu et al., 1978
Oedogonium sp.							
Chlorella fusca	NR	NR	NR	NR	1250	NR	Geyer et al., 1984

TABLE 17 (continued)

Species	Water Concentration (ug/L)	Exposure Duration	Tissue	Tissue Concentration (ppb)	BCF	Elimination $t_{1/2}$	Reference
			SALTWATER				
Eel							
Anguilla anguilla	100	8 days	liver	33,400	334	5.8 days	Holmberg et al., 1972
	100	8 days	muscle	9400	94	NR	Holmberg et al., 1972
	100	8 days	blood	4400	44	NR	Holmberg et al., 1972
Killifish	57-610	168 hours	wholefish	33,000(max)	47-64	4.7 days	Trujillo et al., 1982
Fundulus similis	NR	96 hours	wholefish	NR	30	NR	Schimmel et al., 1978
Sheepshead minnow *Cyprinodon variegatus*	NR	NR	wholefish	NR	5-48	NR	Parrish et al., 1978
Striped mullet *Mugil cephalus*	NR	96 hours	wholefish	NR	38	NR	Schimmel et al., 1978
Carpet shell *Tapes philipinarum*	NR	24 hours	wholefish	NR	~20	NR	Kobayashi et al., 1969
Oyster	2.5	28 days	wholefish	NR	78	<4 days	Schimmel et al., 1978
Crassostrea virginica	25	28 days	wholefish	NR	41	<4 days	Schimmel et al., 1978
Mussel *Mytilus edulis*	NR	NR	wholefish	NR	140-370	NR	Geyer et al., 1982
Grass shrimp *Palaemoneles pugio*	NR	96 hours	wholefish	NR	1.7	NR	Schimmel et al., 1978
Brown shrimp *Penaeus aztecus*	NR	96 hours	wholefish	NR	0.26	NR	Schimmel et al., 1978
Polychaete worm							
Lanice conchilega	NR	NR	wholefish	NR	3830	NR	Ernst, 1979
Neanthes virens	100	2 weeks	wholefish	NR	280	NR	Carr and Neff, 1981

NR = Not reported

Kobayashi and Kishino (1980) and Spehar et al. (1985) also explained why the acute and chronic toxicity of pentachlorophenol increased with decreasing pH. At lower pH, the un-ionized form of pentachlorophenol is favored, and since uptake of pentachlorophenol probably occurs primarily by passive diffusion of the un-ionized form through the gill membrane (Spehar et al., 1985), uptake is also enhanced at low pH. The decreased toxicity of pentachlorophenol at higher pH is apparently caused by fish not accumulating pentachlorophenol to toxic levels. This also explains some of the variability in pentachlorophenol toxicity between studies in which pentachlorophenol is administered in ionized (NaPCP) or un-ionized form.

A study by Niimi and Cho (1983) addressed the relative importance of pentachlorophenol uptake from food vs. water. Rainbow trout were fed diets containing 10 or 40 ppb pentachlorophenol for 110 days. Fish in both groups generally accumulated pentachlorophenol to tissue levels of 1-2 μg/kg. Fish fed diets containing 3 ppm accumulated 40 μg/kg after 50 days, but tissue levels then declined to 20 μg/kg after 110 days. These levels were lower than those accumulated by fish exposed to waterborne pentachlorophenol in a previous study (Niimi and McFadden, 1982), and therefore, the authors concluded that uptake of pentachlorophenol from water would be a more important accumulative pathway in the environment than uptake from food. The authors supported this finding with monitoring data from Lake Ontario fishes that showed similar pentachlorophenol residues in forage species and predators, suggesting that pentachlorophenol bioconcentration through the food chain was not occurring.

Environmental monitoring data for pentachlorophenol residues in aquatic organisms are presented in Table 18. This information and that of Table 17 indicate that the liver is the primary site of pentachlorophenol accumulation in fish. Table 18 also indicates that tissue levels of pentachlorophenol in fish are generally <1 μg/g in unpolluted environments; however, extremely high tissue concentrations (10-500 μg/g) may occur in fish exposed to high levels of pentachlorophenol (Pierce, 1978; Pierce and Victor, 1978). Pierce (1978) and Pierce and Victor (1978) monitored pentachlorophenol in a freshwater lake where accidental release of wood-treating wastes containing pentachlorophenol caused extensive fish kills. They found that pentachlorophenol persisted in water and fish for 6 months following the spills. Concentrations in fishes declined as the concentration in water declined, but it took 6-10 months for fish tissue concentrations to return to normal levels.

Other Relevant Information. Several authors have investigated effects of pentachlorophenol on experimental ecosystems. Schauerte et al. (1982) reported a decline in *Daphnia* populations, blue-green algae and diatoms, and an increase in flagellate protozoans and heterotrophic bacteria in ponds to which 1 mg/ℓ pentachlorophenol was added. Hedtke and Arthur (1985) studied effects of pentachlorophenol on outdoor experimental streams. Snails were affected only at the highest concentration, 432 μg/ℓ, but fish, periphyton and system metabolism occurred at all concentrations tested (48, 144 and 432 μg/ℓ . Differences between the 48 μg/ℓ stream and the control were slight and may not have been caused by pentachlorophenol.

TABLE 18
Pentachlorophenol Residues Detected in Aquatic Biota

Species	Location	Tissue	Tissue Concentration (ng/g)	Reference
		FRESHWATER		
Sucker unspecified	background, Mississippi Lake	muscle	14	Pierce, 1978
	background, Mississippi Lake	liver	380	Pierce, 1978
	after spill, Mississippi Lake	muscle	7,000	Pierce, 1978
	after spill, Mississippi Lake	liver	466,000	Pierce, 1978
	after spill, Mississippi Lake	gill	35,000	Pierce, 1978
Bluegill *Lepomis macrochirus*	background, Mississippi Lake	muscle	4-5	Pierce and Victor, 1978; Pierce, 1978
	background, Mississippi Lake	liver	26-150	Pierce and Victor, 1978; Pierce, 1978
	after spill, Mississippi Lake	muscle	6,400-9,400	Pierce and Victor, 1978; Pierce, 1978
	after spill, Mississippi Lake	liver	130,000	Pierce and Victor, 1978; Pierce, 1978
	experimental ponds after monthly application of 0.2-0.4 mg/L PCP	wholefish	44,000-104,000	Robinson-Wilson et al., 1983
Largemouth bass *Micropterus salmoides*	after spill, Mississippi Lake	gill	48,400	Pierce and Victor, 1978; Pierce, 1978
	after spill, Mississippi Lake	muscle	7,000-17,000	Pierce and Victor, 1978; Pierce, 1978
	after spill, Mississippi Lake	liver	140,000-325,000	Pierce and Victor, 1978; Pierce, 1978
	experimental ponds after monthly application of 0.2-0.4 mg/L PCP	wholefish	7,000-92,000	Robinson-Wilson et al., 1983
Channel catfish *Ictalurus punctatus*	after spill, Mississippi Lake	gill	42,000	Pierce and Victor, 1978; Pierce, 1978
	after spill, Mississippi Lake	muscle	13,000-19,000	Pierce and Victor, 1978; Pierce, 1978
	after spill, Mississippi Lake	liver	214,000	Pierce and Victor, 1978; Pierce, 1978
	experimental ponds after monthly application of 0.2-0.4 mg/L PCP	wholefish	75,000-182,000	Robinson-Wilson et al., 1983
Rock bass *Ambloplites rupestris*	tributary of Lake Erie, 1.7 km below PCP source	wholefish	10	Metcalfe et al., 1984
Crayfish *Oronectes propinquus*	tributary of Lake Erie, 1.7 km below PCP source	wholefish	1	Metcalfe et al., 1984
Tadpole *Rana catesbeiana*	tributary of Lake Erie, 1.7 km below PCP source	wholefish	19	Metcalfe et al., 1984
Leeches *Hirudinea*	tributary of Lake Erie, 1.7 km below PCP source	wholefish	2,671	Metcalfe et al., 1984
	tributary of Lake Erie, 7.4 km below PCP source	wholefish	1-834	Metcalfe et al., 1984

TABLE 18 (continued)

Species	Location	Tissue	Tissue Concentration (ng/g)	Reference
		FRESHWATER (cont.)		
Silver Lake *Merluccius bilinearis*	St. John Estuary, Canada	wholefish	1.75	Zitko et al., 1974
Atlantic salmon *Salmo salar*	St. Croix Estuary, Canada	wholefish	0.54-1.26	Zitko et al., 1974
Cod *Gadus morhua*	St. Croix Estuary, Canada	wholefish	0.82	Zitko et al., 1974
Winter flounder *Pseudopleuronectes americanus*	St. Croix and St. John Estuary, Canada	wholefish	1.77-3.99	Zitko et al., 1974
Sea raven *Hemipterus americanus*	St. Croix Estuary, Canada	wholefish	<0.5	Zitko et al., 1974
Flounder unspecified	San Luis Pass, Texas	wholefish	2.6	Murray et al., 1981
Killifish *Fundulus similis*	San Luis Pass, Texas	wholefish	5.3	Murray et al., 1981
Brown shrimp *Penaeus aztecus*	San Luis Pass, Texas	wholefish	7.5	Murray et al., 1981
Blue crab *Callinectes sapidus*	San Luis Pass, Texas	wholefish	2.8	Murray et al., 1981
Dwarf squid *Lolliguncula brevis*	San Luis Pass, Texas	wholefish	2.9	Murray et al., 1981
Oyster *Crassostrea virginica*	Galveston Bay, Texas	wholefish	3.4-8.3	Murray et al., 1980
Jellyfish unspecified	Gulf of Mexico	wholefish	0.1-1.0	Murray et al., 1980
Sanddab	Los Angeles County, California	liver	5	Gossett et al., 1983
Halibut	Los Angeles County, California	liver	5	Gossett et al., 1983
Scorpionfish	Los Angeles County, California	liver	74	Gossett et al., 1983

TABLE 18 (continued)

Species	Location	Tissue	Tissue Concentration (ng/g)	Reference
		FRESHWATER (cont.)		
Snails *Physa* sp.	tributary of Lake Erie, 7.4 km below PCP source	wholefish	1-9	Metcalfe et al., 1984
Worms *Oligochaeta*	tributary of Lake Erie, 7.4 km below PCP source	wholefish	180-314	Metcalfe et al., 1984
Dragonflies *Anisoptera*	tributary of Lake Erie, 7.4 km below PCP source	wholefish	18-64	Metcalfe et al., 1984
Damselflies *Zygoptera*	tributary of Lake Erie, 7.4 km below PCP source	wholefish	20-34	Metcalfe et al., 1984
Fishflies *Nigronia* sp.	tributary of Lake Erie, 7.4 km below PCP source	wholefish	7-15	Metcalfe et al., 1984
Caddisflies *Hydropsychidae*	tributary of Lake Erie, 7.4 km below PCP source	wholefish	3-6	Metcalfe et al., 1984
Pychopsyche sp.	tributary of Lake Erie, 7.4 km below PCP source	wholefish	15-45	Metcalfe et al., 1984
Craneflies *Tipulidae*	tributary of Lake Erie, 7.4 km below PCP source	wholefish	<1-68	Metcalfe et al., 1984
Water pennies *Psephenidae*	tributary of Lake Erie, 7.4 km below PCP source	wholefish	10-19	Metcalfe et al., 1984
Mayflies *Ephemeroptera*	tributary of Lake Erie, 7.4 km below PCP source	wholefish	12-24	Metcalfe et al., 1984
Pike *Esox* sp.	sites in Finland	wholefish	5.7-8.0	Paasivirta et al., 1980
Roach *Rutilis* sp.	sites in Finland	wholefish	0.9-12.8	Paasivirta et al., 1980
Mussel unspecified	sites in Finland	wholefish	1.7-5.6	Paasivirta et al., 1980
Fishes unspecified	Columbia River, United States, after Mt. St. Helens eruption	wholefish	75-324	Campbell et al., 1982

TABLE 18 (continued)

Species	Location	Tissue	Tissue Concentration (ng/g)	Reference
		FRESHWATER (cont.)		
Croaker	Los Angeles County, California	liver	8	Gossett et al., 1983
Dover sole	Los Angeles County, California	liver	70	Gossett et al., 1983
Crab 0 unspecified	Los Angeles County, California	digestive gland	19	Gossett et al., 1983
Shrimp unspecified	Los Angeles County, California	muscle	5	Gossett et al., 1983
Invertebrates unspecified	Los Angeles County, California	whole	24	Gossett et al., 1983
Polychaete worm Lanice conchilega	estuary, Germany	whole	103-329	Ernst and Weber, 1978
Actinian Sagartia troglodytes	estuary, Germany	whole	2.7-7.0	Ernst and Weber, 1978

Saltwater communities were also affected similarly by sublethal pentachlorophenol concentrations. Tagatz et al. (1977) exposed estuarine benthic communities to 0, 7, 76 or 622 μg/ℓ pentachlorophenol in flowthrough aquaria for 9 weeks. Almost no animals were present at the highest concentration. There were fewer mollusks at 7 μg/ℓ fewer annelids and arthropods at 76 μg/ℓ compared with controls. Total numbers of individuals and species were significantly reduced compared with controls at 76 μg/ℓ not at 7 μg/ℓ . In a similar study, Tagatz et al. (1981) exposed estuarine benthic communities to 0, 1, 13 or 140 μg/ℓ pentachlorophenol for 7 days. Community structure was significantly altered only at the highest concentration. In another study, Tagatz et al. (1978) found that benthic communities were unaffected by 1.8 μg/ℓ , but that 15.8 and 161 μg/ℓ caused a reduction in number of individuals and species. Mollusks were the most sensitive taxonomic group. Cantelmo and Rao (1978) studied effects of pentachlorophenol on mecobenthic nematodes, and found that exposure to 161 or 622 μg/ℓ for 9 weeks resulted in changes in nematode species composition and shifts from epistrate feeders to deposit feeders. Aquaria treated with 1.8, 7, 15.8 and 76 μg/ℓ were similar to controls.

Some of the variability in aquatic toxicology data may be explained by the types of pentachlorophenol used in different studies. Cleveland et al. (1982) compared the toxicity of purified pentachlorophenol and commercial pentachlorophenol in fathead minnow 90-day partial life cycle tests, and found that for purified pentachlorophenol, growth was reduced at concentrations of ≥85 μg/ℓ , and that survival was reduced at 142 μg/ℓ . Commercial pentachlorophenol was much more toxic, with growth reductions occurring at ≥13 μg/ℓ and reduced survival at ≥27 μg/ℓ . These results suggest that impurities present in commercial pentachlorophenol greatly increase its toxicity. Analysis of the commercial pentachlorophenol showed that it contained "relatively large" amounts of hexachlorobenzene, chlorinated phenoxyphenols, chlorinated dibenzodioxins and chlorinated dibenzofurans.

Summary. Acute toxicity data indicated that salmonids are more sensitive to the toxic effects of pentachlorophenol than other fish species, with LC_{50} values of 34-128 μg/ℓ for salmonids and 60-600 μg/ℓ for other species. More recent data showed that carp larvae, bluegills, channel catfish and knifefish also had LC_{50} values <100 μg/ℓ (see Table 10). The most sensitive marine fishes were pinfish larvae, the goby, *Gobius minutus*, and eggs and larvae of the flounder, *Pleuronectes platessa*, all with LC_{50} values <100 μg/ℓ (Adema and Vink, 1981). The most sensitive freshwater invertebrate species were the chironomid, *Chironomous gr. thummi* (Sloof, 1983), and the snail, *Lymnaea luteola* (Gupta et al., 1984). The most sensitive marine invertebrates were the Eastern oyster (Borthwick and Schimmel, 1978), larvae of the crustaceans, *Crangon crangon* and *Palaemon elegans* (Van Dijk et al., 1977), and the copepod, *Pseudodiaptomus coronatus* (Hauch et al., 1980); all had LC_{50} values <200 μg/ℓ .

In chronic toxicity tests, the lowest concentration reported to cause adverse effects was 1.8 μg/ℓ (NaPCP), which inhibited growth of sockeye salmon (Webb and Brett, 1973). The marine species tested displayed similar thresholds for chronic toxicity.

Both acute and chronic toxicity increased at lower pH, probably because a lower pH favors the un-ionized form of pentachlorophenol, which is taken up more readily and is therefore more toxic than ionized pentachlorophenol (Kobayashi and Kishino, 1980; Spehar et al., 1985).

Data concerning the effects of pentachlorophenol on aquatic plants were highly variable; therefore, it was difficult to draw conclusions from these data.

Pentachlorophenol did not appear to bioaccumulate in aquatic organisms to very high concentrations. BCFs for pentachlorophenol were <1000 for most species tested. The highest BCF was 3830 for the polychaete, *Lanice conchilega* (Ernst, 1979). Some species appear to have an inducible pentachlorophenol-detoxification mechanism, as evidenced in several experiments in which pentachlorophenol tissue levels peaked in 4-8 days and declined thereafter despite continued exposure (Pruitt et al., 1977; Trujillo et al., 1982). A study by Niimi and Cho (1983) indicated that uptake of waterborne pentachlorophenol by gills was much greater than uptake from food, indicating that bioconcentration of pentachlorophenol through the food chain is unlikely. Biomonitoring data of Lake Ontario fishes showed that similar pentachlorophenol levels were found in predators and forage species.

Studies with experimental ecosystems have indicated that ecological effects may occur at pentachlorophenol levels as low as those causing chronic toxicity in sensitive species in single-species tests. The lowest concentration that caused adverse effects in these studies was 15.8 μg/ℓ, which caused a reduction in numbers of individuals and species in a marine benthic community (Tagatz et al., 1978).

EXISTING GUIDELINES AND STANDARDS

Human. The ACGIH (1980, 1985) recommends a TLV of 0.5 mg/m^3 for pentachlorophenol, with a STEL of 1.5 mg/m^3 to prevent cardiovascular injury, and warns of the potential contribution that is due to dermal absorption. OSHA reports an occupational standard and permissible exposure limit of 0.5 mg/m^3, with a warning that potentially harmful amounts of pentachlorophenol can be absorbed through the skin (CFR, 1985).

The US. EPA (1985b) reported an ADI of 0.03 mg/kg/day based on a rat chronic oral NOEL of 3 mg/kg/day (Schwetz et al., 1977) and an uncertainty factor of 100. The 1984 WHO ADI of 0.003 mg/kg/day for pentachlorophenol was based on unspecified animal data and an uncertainty factor of 1000 (U.S. EPA, 1985c). The National Research Council (NRC, 1977) derived a drinking water limit of 0.021 mg/ℓ pentachlorophenol based on a NOEL of 3 mg/kg/day from an 8-month and a 90-day oral rat study and an uncertainty factor of 1000.

Aquatic. Although U.S. EPA (1980a) did not recommend a criterion to protect aquatic biota from the effects of pentachlorophenol, it stated that acute and chronic toxicity to freshwater life occurred at concentrations as low as 55 and 3.2 μg/ℓ, respectively,

and would occur at lower concentrations in species more sensitive than those tested. For marine biota, acute and chronic toxicity reportedly occurred at concentrations as low as 53 and 34 μg/ℓ, respectively, and would occur at lower concentrations in species more sensitive than those tested. More recent studies have determined lower thresholds. For freshwater life, acute toxicity occurred at concentrations as low as 9.5 μg/ℓ (96-hour LC_{50} for carp larvae) (Verma et al., 1981). The lowest reported concentration that caused chronic toxicity was 1.8 μg/ℓ, which inhibited growth of sockeye salmon (Webb and Brett, 1973). The lowest reported acutely toxic concentration for marine organisms was 38 μg/ℓ for pinfish larvae (Borthwick and Schimmel, 1978), which U.S. EPA (1980a) reported but apparently did not use in specifying thresholds. The lowest reported chronic toxicity threshold was still 34 μg/ℓ, which inhibited shell deposition in oysters (Schimmel et al., 1978).

Verma et al. (1981) recommended a MATC of 0.5-0.6 μg/ℓ and cited reported safe concentrations of 0.091-5.5 μg/ℓ. No other guidelines and standards for pentachlorophenol could be located in the available literature as cited in the Appendix.

RISK ASSESSMENT

Several toxicity studies (Schwetz et al., 1974; Goldstein et al., 1977; Kimbrough and Linder, 1978; Knudsen et al., 1974; Johnson et al., 1973; Kerkvliet et al., 1982) compared the effects of oral exposure to a commercial pentachlorophenol solution with the effects of oral exposure to a technical or "purified" pentachlorophenol solution. Because the technical and purified pentachlorophenol solutions contained fewer contaminants than the commercial preparations, only the results of the experiments using technical purified solutions were reported in detail. Many of the technical or purified preparations of pentachlorophenol are contaminated with small but measurable amounts of highly toxic substances, such as dibenzodioxins; therefore, in order to assist the process of quantitative risk assessment, a summary of the chemical composition of the various pentachlorophenol solutions used in the studies that impact the risk assessment is presented in Table 19.

Lifetime oral exposure to pentachlorophenol was not carcinogenic in mice (BRL, 1968; Innes et al., 1969) or rats (Schwetz et al., 1977), regardless of the composition of the pentachlorophenol solution tested. Chronic feeding studies are being conducted in mice; the quality assessment of a study of technical pentachlorophenol and the histopathology of a study of Dowicide EC-7 are in progress (NTP, 1985). Although there are a few studies that suggest pentachlorophenol may be slightly mutagenic in *B. subtilis* (Waters et al., 1982; Shirasu, 1976) as evidenced by the rec assay and in mice as evidenced by the coat color spot test (Fahrig et al., 1977), no evidence of mutagenicity was reported in *S. typhimurium* (Anderson et al., 1972; Simmon et al., 1977; Lemma and Ames, 1975; Moriya et al., 1983; Waters et al., 1982; Buselmaier et al., 1973) or in *E. coli* (Simmon et al., 1977; Fahrig, 1974; Moriya et al., 1983; Waters et al., 1982) with or without metabolic activation.

TABLE 19
Selected Contaminants of Technical or Purified Pentachlorophenol[a]

% Pentachlorophenol	Dibenzofurans (ppm)	Dibenzodioxins (ppm)	Reference
90	6.2	22.55	Schwetz et al., 1977[b]
>99	<0.1	<0.1	Kimbrough and Linder, 1978
>99	<0.1	<0.1	Goldstein et al., 1977
NR	NR	282	Knudsen et al., 1974
88-93	NR	31	Johnson et al., 1973
99	NR	NR	Kerkvliet et al., 1982
>98	<1.5	<2	Schwetz et al., 1974
NR	NR	NR	Courtney et al., 1976
85.5	NR	408	Exon and Koller, 1982

[a]In some commercial grade pentachlorophenol, the concentrations of the contaminants were as high as 2600 ppm for dibenzodioxins (Schwetz et al., 1977) and 900 ppm for dibenzofurans (Goldstein et al., 1977).

[b]Test substance used in this study also contained 400 ppm hexachlorobenzene.

NR = Not reported

Three teratogenicity/reproductive toxicity studies (Schwetz et al., 1974, 1977; Courtney et al., 1976) indicate that pentachlorophenol is fetotoxic to rats at dose levels ≥5 mg/kg/day (Schwetz et al., 1974). Increased resorptions, decreased crown-rump length and decreased fetal body weight occurred at ≥30 mg/kg/day, subcutaneous edema and skeletal anomalies occurred at ≥15 mg/kg/day, and delayed ossification occurred at 5 mg/kg/day. A decrease in litter size, though not statistically significant, was reported at the highest dose tested (500 ppm) in another teratogenicity/reproductive toxicity study (Exon and Koller, 1982); the solution tested in this study had a high dibenzodioxin

content (408 ppm). The dietary concentration of 500 ppm is equivalent to 50 mg/kg/day, assuming that young rats in a subchronic study consume a daily amount equal to 10% of their body weight (Mitruka et al., 1976). No adverse fetal or reproductive effects were reported at oral doses ≤3 mg/kg/day (Schwetz et al., 1977; Exon and Koller, 1982).

Adverse effects produced by subchronic and chronic oral exposure to pentachlorophenol occurred primarily in the liver (Kerkvliet et al., 1982; Johnson et al., 1973; Knudsen et al., 1974; Goldstein et al., 1977; Kimbrough and Linder, 1978; Schwetz et al., 1977), the kidney (Johnson et al., 1973; Kimbrough and Linder, 1978; Schwetz et al., 1977) and the immune system (Kerkvliet et al., 1982). Knudsen et al. (1974) reported increased liver weight in the female rats and centrilobular vacuolization in the male rats exposed for 12 weeks to diets containing ≥50 ppm (~5 mg/kg/day, based on food consumption data provided by the investigators) pentachlorophenol, which contained 282 ppm of dibenzodioxins. Increased liver weights were reported in rats ingesting 10 mg/kg/day pentachlorophenol for 90 days (Johnson et al., 1973). In three 8-month studies, no adverse liver effects were reported in mice exposed to diets containing 50 ppm pentachlorophenol (Kerkvliet et al., 1982) or in rats exposed to diets containing ≥20 ppm of pentachlorophenol (Goldstein et al., 1977; Kimbrough and Linder, 1978). Assuming that mice consume a daily amount of food equal to 13% of their body weight, the 50 ppm dietary level is equivalent to 6.5 mg/kg/day. Based on food consumption data provided by Kimbrough and Linder (1978), 20 ppm is equal to 1 mg/kg/day for rats. Furthermore, no adverse hepatic effects were reported in rats ingesting 1, 3 or 5 mg/kg/day pentachlorophenol for 2 years (Schwetz et al., 1977). Increased pigmentation of hepatocytes was reported in 8/27 female rats ingesting 10 mg/kg/day pentachlorophenol for 2 years (Schwetz et al., 1977).

Increased kidney weights unaccompanied by renal histopathology were reported in male rats exposed to 20, 100 or 500 ppm (1, 5 or 29 mg/kg/day based on food consumption data) for 8 months (Kimbrough and Linder, 1978) and in rats (sex unspecified) ingesting 30 mg/kg/day of pentachlorophenol for 90 days (Johnson et al., 1973). Increased pigmentation of the renal tubular epithelial cells was reported in female rats ingesting 10 or 30 mg/kg/day pentachlorophenol for 2 years (Schwetz et al., 1977). Since increased kidney weight in the absence of histopathological changes is not considered an adverse effect, the lowest dose at which adverse renal effects occurred is 10 mg/kg/day pentachlorophenol (Schwetz et al., 1977).

Although decreased immunocompetence was reported in mice exposed to diets containing 50 or 500 ppm (6.5 or 65 mg/kg/day) pentachlorophenol for 34 weeks (Kerkvliet et al., 1982), the decrease was statistically significant only at the higher dose. Therefore, based on the subchronic/chronic toxicity data and disregarding the Knudsen et al. (1974) study because of the high dioxin content of the solution used, the highest NOAEL below which no adverse effects were observed is 5 mg/kg/day pentachlorophenol from the 2-year feeding study using rats (Schwetz et al., 1977). The teratogenicity study of Schwetz et al. (1974) indicated, however, that there was a

significant delay in cranial ossification in pups whose dams were treated by gavage with 5 mg/kg/day pentachlorophenol on days 6-15 of gestation. The next lower NOAEL in the subchronic/chronic studies is 3 mg/kg/day; this dose did not result in fetotoxicity in the dietary reproduction study of Schwetz et al. (1977). In order to protect against possible fetotoxic effects of pentachlorophenol, the ADI of 0.03 mg/kg/day or 2.1 mg/day for a 70 kg human is obtained by dividing the animal NOAEL of 3 mg/kg/day from the 2-year feeding study using rats (Schwetz et al., 1977) by an uncertainty factor of 100 to account for interspecies extrapolation and the range of sensitivity within the human population to xenobiotics.

REPORTABLE QUANTITIES

Reportable Quantity (RQ) Ranking Based on Chronic Toxicity. The effect levels from the chronic/subchronic toxicity studies and teratogenicity/reproductive toxicity studies are listed in Table 20. Fetotoxicity, immunocompetence, hepatotoxicity and renal toxicity are the four areas in which effects occur for consideration in the derivation of the RQ.

Contamination with dibenzofurans and dibenzodioxins, which are known to cause adverse effects when administered in minute amounts, often occurs in purified preparations of pentachlorophenol. The reported purity of pentachlorophenol and the amount of contamination with dibenzofurans and dibenzodioxins in each of the studies under consideration are listed in Table 19.

Results from four teratogenicity/reproductive toxicity studies (Schwetz et al., 1974, 1977; Courtney et al., 1976; Exon and Koller, 1982) indicate that pentachlorophenol is fetotoxic at dose levels $\geq$5 mg/kg/day. Evaluation of fetal anomalies and skeletal variations, however, was performed only in the Schwetz et al. (1974, 1977) studies. In the Schwetz et al. (1977) study, the rats were treated with dietary concentrations of 0, 3 or 30 mg/kg/day pentachlorophenol for 62 days before mating, and continuing throughout mating, gestation and lactation. At 30 mg/kg/day, but not at 3 mg/kg/day, weanling rats had skeletal variations. In the Schwetz et al. (1974) study, the rats were treated by gavage with 0, 5, 15, 30 or 50 mg/kg/day pentachlorophenol on days 6-15 of gestation. Fetal anomalies, subcutaneous edema and dilated ureters occurred at 30 mg/kg/day, and skeletal variations, including lumbar spurs, unfused vertebral centers, extra ribs, delayed cranial, vertebral and sternal ossification occurred at $\geq$15 mg/kg/day. The most consistently occurring skeletal change, delayed cranial ossification, was observed at the lowest dose, 5 mg/kg/day. Thus, the MED of 59.8 mg/day for adverse fetal effects was derived by multiplying the animal dose of 5 mg/kg/day from the Schwetz et al. (1974) study by the cube root of the rat to the reference human body weight ratio (assumed to be 0.35 kg for a rat and 70 kg for a human) and by 70 kg. As indicated in Table 21, the RV_d corresponding to a human MED of 59.8 mg/day is 2.8; the appropriate RV_e for fetotoxicity, as evidenced by delayed ossification in the absence of maternal toxic effects is 8. A CS of 22.4 is obtained by multiplying the RV_d by the RV_e, which corresponds to an RQ of 100.

TABLE 20
Oral Toxicity Summary for Pentachlorophenol

Species/ Strain	Sex	No. at Start	Average Weight (kg)	Vehicle/ Physical State	Purity	Exposure	Transformed Animal Dose* (mg/kg/day)	Response	Reference
Rat/ Sprague-Dawley	M/F	27/27	0.35 (assumed)	dietary	90%	30 mg/kg/day for 22-24 months	30	Increased SGPT levels in both males and females; more than half the females, but only 1/27 males, had brown pigment present in hepatocytes; 19/27 females had brown pigment in renal tubular epithelial cells	Schwetz et al., 1977
Rat/ Sprague-Dawley	M/F	27/27	0.35 (assumed)	dietary	90%	10 mg/kg/day for 24 months	10	Brown pigment reported in hepatocytes of 8/27 females and in renal tubular epithelial cells of 7/27 females; no effects in males	Schwetz et al., 1977
Rat/ Sherman	M/F	10/10	0.40M/0.25F (experimental)	dietary	>99%	500 ppm for 8 months	29M/32F	Decreased body weight in both males and females; centrilobular hepatocyte enlargement in eight males and five females; increased kidney weight in males	Kimbrough and Linder, 1978
Rat/ Sherman	M/F	10/10	0.45M/0.27SF (experimental)	dietary	>99%	100 ppm for 8 months	5M/6F	Increased kidney weight in males without histopathology	Kimbrough and Linder, 1978
Rat/ Sherman	M/F	10/10	0.45M/0.27SF (experimental)	dietary	>99%	20 ppm for 8 months	1M/1.1F	Increased kidney weight in males without histopathology	Kimbrough and Linder, 1978
Rat/ Sherman	F	6	0.35 (assumed)	dietary	>99%	500 ppm for 8 months	25 (assumed)	Significantly decreased body weight; significant increase in hepatic p-nitrophenol glucuronyl transferase	Goldstein et al., 1977
Rat/ Wistar	M/F	10/10	0.204/0.107 (experimental)	dietary	NR	200 ppm for 12 weeks	18.3M/21.7F	In females, increased liver weight and hepatic aryl hydrolase levels, increased serum alkaline phosphatase levels; in males, increased hepatic centrilobular vacuolization (5/10) and aryl hydrolase levels, decreased hemoglobin levels and red blood cell counts and significantly increased serum glucose levels	Knudsen et al., 1974

TABLE 20 (continued)

Species/ Strain	Sex	No. at Start	Average Weight (kg)	Vehicle/ Physical State	Purity	Exposure	Transformed Animal Dose* (mg/kg/day)	Response	Reference
Rat/ Wistar	M/F	10/10	0.204/0.107 (experimental)	dietary	NR	50 ppm for 12 weeks	4.34M/5.47F	In males, increased incidence of hepatic centrilobular vacuolization (4/10); decreased white blood cell count and decreased hepatic glucose-6-phosphatase; no adverse effects in females	Knudsen et al., 1974
Rat/ Wistar	M/F	10/10	0.204/0.107 (experimental)	dietary	NR	25 ppm for 12 weeks	2.12M/2.69F	In females, significantly decreased body weight and significantly increased serum urea and alkaline phosphatase levels; no adverse effects in males	Knudsen et al., 1974
Rat/ Sprague-Dawley	NR	NR	0.35 (assumed)	dietary	88-93%	30 mg/kg/day for 90 days	30	Increased liver and kidney weight without associated histopathology	Johnson et al., 1973
Rat/ Sprague-Dawley	NR	NR	0.35 (assumed)	dietary	88-93%	10 mg/kg/day for 90 days	10	Increased liver weight without associated histopathology	Johnson et al., 1973
Mouse/ C57B/ 6(B6)	M	10	0.03 (assumed)	dietary	>99%	500 ppm for 34 weeks	65 (assumed)	Increased immunosuppression as evidenced by splenic tumors in 4/9 mice (controls 0/13) and mild multifocal necrosis associated with hepatocyte swelling and vacuolization	Kerkvliet et al., 1982
Mouse C57BL/ 6(B6)	M	10	0.03 (assumed)	dietary	>99%	50 ppm for 34 weeks	6.5 (assumed)	Increased immunosuppression as evidenced by splenic tumors in 2/9 mice (controls 0/13)	Kerkvliet et al., 1982
Rat/ Sprague-Dawley	F	20	0.35 (assumed)	gavage in corn oil	>98%	50 mg/kg/day on days 6-15 of gestation	50	100% resorption	Schwetz et al., 1974
Rat/ Sprague-Dawley	F	20	0.35 (assumed)	gavage in corn oil	>98%	30 mg/kg/day on days 6-15 of gestation	30	97.5% resorption; decreased fetal birth weight and crown-rump length; increased male: female pup ratio; increased incidence of subcutaneous edema and skeletal anomalies	Schwetz et al., 1974

TABLE 20 (continued)

Species/ Strain	Sex	No. at Start	Average Weight (kg)	Vehicle/ Physical State	Purity	Exposure	Transformed Animal Dose* (mg/kg/day)	Response	Reference
Rat/ Sprague-Dawley	F	20	0.35 (assumed)	gavage in corn oil	>98%	15 mg/kg/day on days 6-15 of gestation	15	Increased incidence of subcutaneous edema, extra ribs, lumbar spurs and delayed cranial ossification	Schwetz et al., 1974
Rat/ Sprague-Dawley	F	20	0.35 (assumed)	gavage in corn oil	>98%	5 mg/kg/day on days 6-15 of gestation	5	Increased incidence of delayed cranial ossification	Schwetz et al., 1974
Rat/C-D	F	7	0.35 (assumed)	gavage in corn oil: acetone (9:1)	NR	75 mg/kg/day on days 7-18 of gestation	75	Decreased fetal weight	Courtney et al., 1976
Rat/ Sprague-Dawley	F	12-20	0.35 (assumed)	dietary	85.5%	500 ppm in food from 21 days post-partum through mating, gestation and lactation	50 (assumed)	Although not statistically significant, decreased litter size	Exon and Koller, 1982
Rat/ Sprague-Dawley	M/F	10/20	0.35 (assumed)	dietary	90%	30 mg/kg/day in the food for 62 days and continued through mating, gestation and lactation	30	Decreased maternal weight at 21 days post-partum; decreased pup weight through lactation; decreased live-born pups; decreased survival rate of pups	Schwetz et al., 1977

*In dietary studies, the transformed dose was supplied by the investigators or was calculated from food consumption data provided by the investigator, unless stated to be assumed. The assumptions in calculating the dose were that rats and mice in a chronic study consume a daily amount of food equal to 5 and 13%, respectively of their body weight (U.S. EPA, 1980b), and that young growing rats in the subchronic study by Exon and Koller (1982) consume a daily amount of food equal to 10% of their body weight (Mitruka et al., 1976).

An RQ of 100 is also derived for the immunological effect reported by Kerkvliet et al. (1982). Transplantation of virus transformed cells into the subcutaneous tissue of mice exposed to 0, 50 ppm (6.5 mg/kg/day) or 500 ppm (65 mg/kg/day) of purified (≥99%) pentachlorophenol in the diet for 34 weeks resulted in splenic tumors in 0/13 controls, 2/9 low-dose mice and 4/9 high-dose mice. Although the increased incidence of splenic tumors was statistically significant only at the high-dose level, reduced immunocompetence apparently occurred at the low-dose level as well. Therefore, the MED of 34.3 mg/day for adverse immunological effects was calculated by multiplying the animal dose of 6.5 mg/kg/day by the cube root of the animal to human body weight ratio (assumed to be 0.03 kg for a mouse and 70 kg for a human) and by the reference human body weight. As indicated in Table 21, the appropriate RV_d for a human MED and 34.4 mg/day is 3.2; the appropriate RV_e for reduced immunocompetence is 7. Multiplying the RV_d by the RV_e results in a CS of 22.4, which corresponds to an RQ of 100.

Adverse liver effects, such as increased liver weight and centrilobular vacuolization, occurred in the chronic/subchronic toxicity studies (Schwetz et al., 1977; Kimbrough and Linder, 1978; Goldstein et al., 1977; Knudsen et al., 1974; Johnson et al., 1973; Kerkvliet et al., 1982). The only study in which adverse liver effects occurred at dose levels <10 mg/kg/day is the 12-week Knudsen et al. (1974) study, in which increased liver weight was reported in female rats at 5.47 mg/kg/day pentachlorophenol and centrilobular vacuolization was reported in male rats at 4.34 mg/kg/day pentachlorophenol. Use of the Knudsen et al. (1974) study to derive an RQ for pentachlorophenol is not appropriate, however, because the pentachlorophenol used in testing contained 282 ppm dioxins, ≥10 times the content of dioxins contained in the test solution used by Schwetz et al. (1977) in a 2-year toxicity study in which no adverse liver effects were observed in rats receiving 1, 3 or 5 mg/kg/day pentachlorophenol. As indicated in Table 19, the dioxin contents were 31 and 22.55 ppm in the Johnson et al. (1973) and Schwetz et al. (1977) studies, respectively. In the 90-day Johnson et al. (1973) study, increased liver weight but no histopathology was observed in the rats (sex unspecified) treated with 10 mg/kg/day pentachlorophenol. In the Schwetz et al. (1977) study, rats had increased pigmentation of hepatocytes at 10 mg/kg/day for 2 years. Therefore, the MED of 119.7 mg/day for adverse liver effects was derived by multiplying the animal dose of 10 mg/kg/day from the Johnson et al. (1973) and Schwetz et al. (1977) studies by the cube root of the animal to human body weight ratio (assumed to be 0.35 kg for a rat and 70 kg for a human) and by the reference human body weight. As indicated in Table 21, the appropriate RV_d for a human MED of 119.7 mg/day is 2.4; the appropriate RV_e for increased liver weight and/or increased pigmentation of hepatocytes is 5. Multiplying the RV_d by the RV_e results in a CS of 12, which corresponds to an RQ of 1000.

Increased pigmentation of the tubular epithelial cells was reported in female rats, but not males receiving 10 or 30 mg/kg/day dietary pentachlorophenol (Schwetz et al., 1977). Increased kidney weight was reported in male rats, but not in females, at all

TABLE 21
Composite Scores for Oral Toxicity of Pentachlorophenol

Species	Animal Dose (mg/kg/day)	Chronic Human MED (mg/day)	RV_d	Effect	RV_e	CS	RQ	Reference
Rat	5	59.8	2.8	Fetotoxicity as evidenced by delayed ossification; no maternotoxicity	8	22.4	100	Schwetz et al., 1974
Mice	6.5	34.3	3.2	Immunosuppression as evidenced by increased incidence of splenic tumors after implantation of virus transformed tumor cells	7	22.4	100	Kerkvliet et al., 1982
Rats	10	119.7	2.4	Increased liver weight without any histopathology; increased incidence of pigmented hepatocytes	5	12	1000	Johnson et al., 1973; Schwetz et al., 1977
Rats	1	13.0	8	Increased kidney weights in males without any histopathology	4	15.2	1000	Kimbrough and Linder, 1978

dose levels (1, 5 and 29 mg/kg/day) in the 8-month study by Kimbrough and Linder (1978). The test solution of pentachlorophenol used by Kimbrough and Linder (1978) was ≥99% pure and the concentration of dibenzofurans and dibenzodioxins was <0.1 ppm, the lower limit of detection (see Table 19). Even though the increased kidney weight was not dose-dependent and the 2-year study by Schwetz et al. (1977) did not report increased kidney weight in male or female rats at any dose level tested (1, 3, 5, 10 or 30 mg/kg/day of pentachlorophenol), the animal dose of 1 mg/kg/day was used to derive an RQ. Therefore, the MED of 13 mg/day for adverse renal effects was derived by multiplying the animal dose of 1 mg/kg/day from the Kimbrough and Linder (1978) study by the cube root of the animal to human body weight ratio (reported to be 0.45 for male rats and 70 kg for humans) and by 70 kg. As indicated in Table 21, the appropriate RV_d for a human MED of 13 mg/day is 3.8; the appropriate RV_e for increased kidney weight is 4. Multiplying the RV_e by the RV_d results in a CS of 15.2, which corresponds to an RQ of 1000.

As indicated by Table 21 and reported in Table 22, the most appropriate RQ for pentachlorophenol from the available data is 100, based on the CS of 22.4 obtained from the fetotoxic effects reported by Schwetz et al. (1974). An RQ of 100 is supported by the decreased immunocompetence reported in mice exposed to dietary pentachlorophenol (Kerkvliet et al., 1982).

Weight of Evidence of Potency Factor ($F = 1/ED_{10}$) for Carcinogenicity. An increased incidence of cancer was not observed after oral (BRL, 1968; Innes et al., 1969; Schwetz et al., 1977), subcutaneous (BRL, 1968; Catalina et al., 1981) or dermal (Boutwell and Bosch, 1959) exposure to pentachlorophenol. The quality assessment of a chronic feeding study of technical pentachlorophenol in mice and the histological analyses of a chronic feeding of Dowicide EC-7 in mice are in progress (NTP, 1985).

TABLE 22
Pentachlorophenol
Minimum Effective Dose (MED) and Reportable Quantity (RQ)

Route:	oral
Dose*:	59.8 mg/day
Effect:	fetotoxicity as evidenced by delayed ossification
Reference:	Schwetz et al., 1974
RV_d:	2.8
RV_e:	8
Composite Score:	22.4
RQ:	100

*Equivalent human dose

Pentachlorophenol was not mutagenic in *S. typhimurium, E. coli* (Anderson et al., 1972; Simmon et al., 1977; Lemma and Ames, 1975; Moriya et al., 1983; Waters et al., 1982; Buselmaier et al., 1973; Fahrig, 1974) or *Drosophila* (Vogel and Chandler,

1974). There is some evidence that pentachlorophenol may be weakly mutagenic in *B. subtilis* (Waters et al., 1982; Shirasu, 1976) and in mice as evidenced by the coat color spot test (Fahrig et al., 1977). Therefore, according to IARC (1979) and after considering more recent evidence (Schwetz et al., 1977; Catalina et al., 1981), pentachlorophenol belongs in IARC Group 3, that is, it cannot be classified as to its carcinogenicity for humans. According to U.S. EPA (1984b), pentachlorophenol belongs in CAG Group D, that is, there is no evidence of carcinogenicity for humans. These classifications may change pending the result of the NTP studies.

DISCLAIMER

This report is an external draft for review purposes only and does not constitute Agency policy. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

REFERENCES

ACGIH (American Conference of Governmental Industrial Hygienists). (1980). Documentation of the Threshold Limit Values, 4th ed., p. 323. Cincinnati, OH.

ACGIH. (1985). TLVs: Threshold Limit Values for Chemical Substances and Physical Agents in the Workroom Environment with Intended Changes for 1984–1985, p. 26. Cincinnati, OH.

ADEMA, D.M.M. (1978). *Daphnia magna* as a test animal in acute and chronic toxicity tests. Hydrobiologia **59**:125–134.

ADEMA, D.M.M. and VINK, I.G.J. (1981). A comparative study of the toxicity of 1,1,2-trichloroethane, dieldrin, pentachlorophenol and 3,4-dichloroaniline for marine and fresh water organisms. Chemosphere **10**:533–554.

AHLBORG, U.H. and THUNBERG, T.M. (1980). Chlorinated phenols: Occurrence, toxicity, metabolism and environmental impact. CRC Crit. Rev. Toxicol. **7**:1–35.

AHLBORG, U.G., LINDGREN, J.E. and MERCIER, M. (1974). Metabolism of pentachlorophenol. Arch. Toxicol. **32**:271–281.

ALABASTER, J.S. (1957). Toxicity of weedkilling, algicides and fungicides to trout. In: Proc. 4th Br. Weed Cont. Conf., Brighton, p. 84. (Cited in U.S. EPA, 1980a).

ALEXANDER, M. and ALEEM, M.I.H. (1961). Effect of chemical structure on microbial decomposition of aromatic herbicides. J. Agric. Food Chem. **9**:44–47.

ALEXANDER, H.C., BODNER, K.M. and MAYES, M.A. (1983). Evaluating the "OECD Fish Prolonged Toxicity Study at Least 14 Days." Chemosphere **12**:415–423.

ANDERSON, K.J., LEIGHTY, E.G. and TAKAHASHI, M.T. (1972). Evaluation of herbicides for possible mutagenic properties. J. Agric. Food Chem. **20**:649–656.

BAKER, M.D. and MAYFIELD, C.I. (1980). Microbial and non-biological decomposition of chlorophenols and phenols in soil. Water Soil Pollut. **13**:411.

BAKER, M.D., MAYFIELD, C.I. and INNISS, W.E. (1980). Degradation of chlorophenol in soil, sediment and water at low temperatures. Water Res. **14**:1765–1771.

BANERJEE, S., HOWARD, A.M., ROSENBERG, A.E., DOMBROWSKI, A.E., SIKKA, H. and TULLIS, D.L. (1984). Development of a general kinetic model for biodegradation and its application to chlorophenols and related compounds. Environ. Sci. Technol. **18**:416–422.

BEVENUE, A., OGATA, J.N. and HYLIN, J.W. (1972). Organochlorine pesticides in rainwater, Oahu, Hawaii, 1971–1972. Bull. Environ. Contam. Toxicol. **8**:238–241.

BORTHWICK, P.W. and SCHIMMEL, S.C. (1978). Toxicity of pentachlorophenol and related compounds to early life stages of selected estuarine animals. In: Chemistry, Pharmacology and Environmental Toxicology, K. R. Rao, ed. Plenum Press: New York.

BOURQUIN, A.W. (1984). Biodegradation in the estuarine-marine environments and the genetically altered microbe. ERL, U.S. EPA, Gulf Breeze, FL. EPA 600/D-84-051. NTIS PB84-151315.

BOUTWELL, R.K. and BOSCH, D.K. (1959). Tumor-promoting action of phenol and related compounds for mouse skin. Cancer Res. **19**:413–424.

BRASS, H.J., FEIGE, M.A., HALLORAN, T., MELLO, J.W., MUNCH, D. and THOMAS, R.B. (1977). The national organic monitoring survey: Samplings and analyses for purgeable organic compounds and drinking water quality. Enhancement Source Prot., pp. 393–416.

BRAUN, W.H. and SAUERHOFF, M.W. (1976). The pharmacokinetic profile of pentachlorophenol in monkeys. Toxicol. Appl. Pharmacol. **38**:525–533.

BRAUN, W.H., YOUNG, J.D., BLAU, G.E. and GEHRING, P.J. (1977). The pharmacokinetics and metabolism of pentachlorophenol in rats. Toxicol. Appl. Pharmacol. **41**:395–406.

BRAUN, W.H., BLAU, G.E. and CHENOWETH, M.B. (1978). The metabolism/pharmacokinetics of pentachlorophenol in man, and a comparison with the rat and monkey model. Toxicol. Appl. Pharmacol. **45**:278.

BRL (Bionetics Research Laboratories). (1968). Evaluation of the Carcinogenic, Teratogenic and Mutagenic Activities of Selected Pesticides and Industrial Chemicals. Vol. 1. Carcinogenic Study, Publ. No. NCI-DCCP-CG-1973-1-1. Prepared by BRL, Bethesda, MD, under Contract Nos. PH 43-64-57 and PH 43-67-735. National Cancer Institute, Bethesda, MD, NTIS PB-223-159.

BUSELMAIER, W., ROHRBORN, G. and PROPPING, P. (1973). Comparative investigations of the mutagenicity of pesticides in mammalian test systems. Mutat. Res. **21**:25. (Cited in U.S. EPA, 1984a).

BUTTE, W., DENKER, J., KIRSCH, M. and HOEPNER, T. (1985). Pentachlorophenol and tetrachlorophenols in Wadden sediment and clams (*Mya arenaria*) of the Jadebusen after a 14-year period of wastewater discharge containing pentachlorophenol. Environ. Pollut., Ser. B **9**:29–39.

CALLAHAN, M.A., SLIMAK, M.W., GABEL, N.W. et al. (1979a). Water-Related Environmental Fate of 129 Priority Pollutants. Volume 11. U.S. EPA, Washington, DC. EPA 440/4-7-0-029B.

CANTELMO, F.R. and RAO, K.R. (1978). Effect of pentachlorophenol on the meiobenthic nematodes in an experimental system. In: Chemistry, Pharmacology, and Environmental Toxicology. Environmental Science Research Series, Vol. 12, K.R. Rao, ed., pp. 165–174. Plenum Press, New York.

CASARETT, B., YAUGER, A., JR., WHALEN, W.L. and SA, L.J. (1969). Observations on pentachlorophenol in human blood and urine. Am. Ind. Hyg. Assoc. J. **30**:360–366.

CATALINA, P., CHAMOUX, A., CATILINA, M.J. and CHAMPEIX, J. (1981). Study of the pathogenic properties of substances used as wood protectives: Pentachlorophenol. Arch. Mal. Prof. Med. Trav. Secur. Soc. **42**:334–337. (Fre.) (Cited in the U.S. EPA, 1984a).

CAUTREELS, W., VANCAUMENBERGE, K. and GUZMAN, L.A. (1977). Comparison between the organic fraction of suspended water at a background and an urban station. Sci. Total Environ. **8**:79–88.

CESSNA, A.J. and GROVER, R. (1978). Spectroscopic determination of dissociation constants of selected acidic herbicides. J. Agric. Food Chem. **26**:289–292.

CFR (Code of Federal Regulations). (1985). OSHA (Occupational Safety and Health Administration). Occupational Standards and Permissible Exposure Limits. **29** CFR 1910.1000.

CHU, J. and KIRSCH, J. (1973). Utilization of halophenols by a pentachlorophenol metabolizing bacterium. Dev. Ind. Microbiol. **14**:264–273.

CLEVELAND, L., BUCKLER, D.R., MAYER, F.L. and BRANSON, D.R. (1982). Toxicity of three preparations of pentachlorophenol to fathead minnows—A comparative study. Environ. Toxicol. Chem. **1**:205–212.

CMR (Chemical Marketing Reporter). (1980). Chemical profile—Pentachlorophenol. October 20, 1980.

COLE, R.H., FREDERICK, R.E., HEALY, R.P. and ROLAN, R.G. (1984). Preliminary findings of the priority pollutant monitoring project of the nationwide urban runoff program. J. Water Pollut. Control Fed. **56**:898–908.

COOK, W.L., FIEDLER, D. and BOURQUIN, A.W. (1980). Succession of microfungi in estuarine microcosms perturbed by carbaryl, methyl parathion and pentachlorophenol. Bot. Mar. **23**:129–131.

COURTNEY, K.D., COPELAND, M.F. and ROBBINS, A. (1976). The effects of pentachloronitrobenzene, hexachlorobenzene, and related compounds on fetal development. Toxicol. Appl. Pharmacol. **35**:239–256.

CROSBY, D.G., BEYNON, K.I., GREVE, P.A., KORTE, F., STILL, G.G. and VONK, J.W. (1981). Environmental chemistry of pentachlorophenol. Pure Appl. Chem. **53**:1051–1080.

CROSSLAND, N.O. and WOLFF, C.J.M. (1985). Fate and biological effects of pentachlorophenol in outdoor ponds. Environ. Toxicol. Chem. **4**:73–86.

CSERJESI, A.J. and JOHNSON, E.L. (1972). Methylation of pentachlorophenol by *Trichoderma vergatum*. Can. J. Microbiol. **18**:45–59.

DALELA, R.C., RANI, S., RANI, S. and VERMA, S.R. (1980). Influence of pH on the toxicity of phenol and its two derivatives, pentachlorophenol and dinitrophenol, to some fresh water teleosts. Acta Hydrochim. Hydrobiol. **8**:623–629.

DANNENBERG, R. (1984). Experiences with a freshwater hydroid test. Z. Wasser Abwasser Forsch. **17**:16–19. CA **101**(1):1917v.

DAVIS, H.C. and HIDU, H. (1969). Effects of pesticides on embryonic development of clams and oysters and on survival and growth of the larvae. Fish. Bull. **67**:393–404.

DELAUNE, R.D., GAMBRELL, R.P. and REDDY, K.S. (1983). Fate of pentachlorophenol in estuarine sediment. Environ. Pollut. (Ser. B) **6**:297–308.

DOBBS, A.J. and WILLIAMS, N. (1983). Indoor air pollution from pesticides used in wood remedial treatments. Environ. Pollut. (Ser. B) **6**:271–296.

DOUGHERTY, R.C. (1978). Human exposure to pentachlorophenol. In: Pentachlorophenol: Chemistry, Pharmacology and Environmental Toxicology, K.R. Rao, ed., pp. 351–361. Pergamon Press, New York.

DRAGUN, J. and HELLING, C.S. (1981). Evaluation of molecular modeling techniques to estimate the mobility of organic chemicals in soils: II. Water solubility and the molecular fragment mobility coefficient. In: Proc. 7th Ann. Res. Land Disposal: Hazardous Waste, pp. 58–70. U.S. EPA, Washington, DC.

DRAHANOVSKY, J. and VACEK, Z. (1971). Dissociation constants and column chromatography of chlorinated phenols. Collect. Czech. Chem. Common. **36**:3431–3440.

DUGGAN, R.E. and CORNELIUSSEN. (1972). Dietary intake of pesticide chemicals in the United States, III. June 1968–April 1970. Pest. Monit. J. **5**:331–341.

DUGGAN, R.E., CORNELIUSSEN, P.E., DUGGAN, M.B., MCMAHON, B.M. and MARTIN, R.J. (1983). Pesticide residue levels in foods in the United States from July 1, 1969 to June 30, 1976. Food and Drug Admin. Div. Chem. Technol., Washington, DC.

DUNCAN, C.G. and DEVERALL, F.J. (1964). Degradation of wood preservatives by fungi. Appl. Microbiol. **12**:57–62.

ECETOC (European Chemical Industry, Ecology and Toxicology Center). (1984). The phototransformation of chemicals in water: Results of a ring-test. Report No. 12. ECETOX, Brussels.

EDER, G. and WEBER, K. (1980). Chlorinated phenols in sediments and suspended matter of the Wester estuary. Chemosphere **9**:111–118.

EDGEHILL, R.U. (1982). Microbial treatment of wastewater and soil to remove pentachlorophenol. Diss. Abstr. Int. B **42**:4871.

EDGEHILL, R.U. and FINN, R.K. (1983). Microbial treatment of soil to remove pentachlorophenol. Appl. Environ. Microbiol. **45**:1122–1125.

ERNST, W. (1979). Factors affecting the evaluation of chemicals in laboratory experiments using marine organisms. Ecotoxicol. Environ. Saf. **3**:90–98. (Cited in U.S. EPA, 1980a).

ERNST, W. and WEBER, K. (1978). Chlorinated phenols in selected estuarine bottom fauna. Chemosphere **7**:867–872.

EXON, J.H. and KOLLER, L.D. (1982). Effects of transplacental exposure to chlorinated phenols. Environ. Health Perspect. **46**:137–140.

FAHRIG, R. (1974). Comparative mutagenicity studies with pesticides. IARC Sci. Publ. **10**:161–181.

FAHRIG, R., NILSSON, C.A. and RAPPE, C. (1977). Genetic activity of chlorophenols and chlorophenol impurities. In: Pentachlorophenol: Chemistry, Pharmacology and Environmental Toxicology, K.R. Rao, ed., pp. 325–338. Plenum Press, New York.

FOX, ME. and JOSHI, S.R. (1984). The fate of pentachlorophenol in the Bay of Quinte, Lake Ontario. J. Great Lakes Res. **10**:190–196.

FREITER, E.R. (1978). Chlorophenols. In: Kirk-Othmer Encyclopedia of Chemical Technology, Vol. 5, 3rd ed., M. Grayson and D. Eckroth, eds., pp. 864–872. John Wiley, New York.

GAEB, S., NITZ, S., PARLAR, H. and KORTE, F. (1975). Photomineralization of certain aromatic xenobiotics. Chemosphere **4**:251–256.

GEBEFUGI, I and KORTE, F. (1983). Pentachlorophenol contamination of human milk samples. Chemosphere **12**:1055–1060.

GLICKMAN, A.H., STATHAM, C.N., WU, A. and LECH, J.J. (1977). Studies on the uptake, metabolism, and disposition of pentachlorophenol and pentachloroanisole in rainbow trout. Toxicol. Appl. Pharmacol. **41**:649–658.

GLUTH, G., FREITAG, D., HANKE, W. and KORTE, F. (1985). Accumulation of pollutants in fish. Comp. Biochem. Physiol., C: Comp. Pharmacol. Toxicol. **81C**(2):273–277.

GOLDSTEIN, J.A., FRIESEN, M., LINDER, R.E. et al. (1977). Effects of pentachlorophenol on hepatic drug-metabolizing enzymes and porphyria related to contamination with chlorinated dibenzo-p-dioxins and dibenzofurans. Biochem. Pharmacol. **26**:1549–1557.

GOSSETT, R.W., BROWN, D.A. and YOUNG, D.R. (1983). Predicting the bioaccumulation of organic compounds in marine organisms using octanol/water partition coefficients. Mar. Pollut. Biol. **14**:387–392.

GUPTA, P.K., RAO, P.S. and MUJUMDAR, V.S. (1984). Studies on the toxicity of pentachlorophenol, sodium pentachlorophenate and phenol to a pulmonate snail *Lymnaea luteola* (Lamarck). Arch. Hydrobiol. **102**:123–130. CA **102**(11):90935k.

HANSCH, C. and LEO, A.J. (1985). MEDCHEM Project. Promona College, Claremont, CA.

HAUCH, R.G., NORRIS, D.R. and PIERCE, R.H. (1980). Acute and chronic toxicity of sodium pentachlorophenate to the copepod, *Pseudodiaptomus coronatus*. Bull. Environ. Contam. Toxicol. **25**:562–568.

HEDTKE, S.F. and ARTHUR, J.W. (1985). Evaluation of a Site-Specific Criterion Using Outdoor Experimental Streams. EPA 600/D-85/041, p. 25. (Taken from Abstract from DIALOG File 6-NTIS).

HELLING, C.S., DENNISON, D.G. and KAUFMAN, D.D. (1974). Fungicide movement in soils. Phytopathology **65**:1091–1100.

HICKMAN, G.T. and NOVAK, J.T. (1984). Acclimation of activated sludge to pentachlorophenol. J. Water Pollut. Control Fed. **56**:364–369.

HINKLE, D.K. (1973). Fetotoxic effects of PCP in the golden syrian hamster. Unpublished. Given at the 12th Annual Meeting of the Society of Toxicology. (Cited in U.S. EPA, 1982).

HOLCOMBE, G.W., PHIPPS, G.L. and FIANDT, J.T. (1982). Effects of phenol, 2,4-dimethylphenol, 2,4-dichlorophenol, and pentachlorophenol on embryo, larval, and early-juvenile fathead minnows (*Pimephales promelas*). Arch. Environ. Contam. Toxicol. **11**:73–78.

HUANG, J.C. and GLOYNA, E.F. (1968). Effect of inorganic compounds on photosynthetic oxygenation I. Chlorophyll destruction and suppression of photosynthetic oxygen production. Water Res. **2**:347–366.

IARC (International Agency for Research on Cancer). (1979). IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Pentachlorophenol. Some Halogenated Hydrocarbons, pp. 309–325. Vol. 20, WHO, IARC, Lyon, France.

IDE, A., NIKI, Y., SAKAMOTO, F. and WATANABE, I. (1972). Decomposition of pentachlorophenol in paddy soil. Agric. Biol. Chem. **36**:1937–1944.

INGOLS, R.S., GAFFNEY, P.E. and STEVENSON, P.C. (1966). Biological activity of halophenols. J. Water Pollut. Control Fed. **38**:629–635.

INNES, J.R.M. et al. (1969). Bioassay of pesticides and industrial chemicals for tumorigenicity in mice. A preliminary note. J. Natl. Cancer Inst. **42**:1101.

JAKOBSON, I. and YLLNER, S. (1971). Metabolism of [^{14}C]-pentachlorophenol in the mouse. Acta Pharmacol. Toxicol. **29**:513–524.

JOHNSON, W.W. and FINLEY, M.T. (1980). Handbook of acute toxicity of chemicals to fish and aquatic invertebrates. Resource Publ. 137. U.S. Dept. Interior, Fish and Wildlife Service, Washington, DC.

JOHNSON, R.L., GEHRING, P.J., KOCIBA, R.J. and SCHWETZ, B.A. (1973). Chlorinated dibenzodioxins and pentachlorophenol. Environ. Health Perspect., Exp. Issue No. **5**:171.

KARNS, J.S., KILBANE, J.J., DUTTAGUPTA, S. and CHAKRABARTY, A.M. (1983). Metabolism of halophenols by 2,4,5-trichlorophenoxyacetic acid-degrading *Pseudomonas cepacia*. Appl. Environ. Microbiol. **46**:1176–1181.

KENAGA, E.E. (1980). Predicted bioconcentration factors and soil sorption coefficients of pesticides and other chemicals. Ecotox. Environ. Saf. **4**:26–38.

KERKVLIET, N.I., BAECHER-STEPPAN, L. and SCHMITZ, J.A. (1982). Immunotoxicity of pentachlorophenol (PCP): Increased susceptibility to tumor growth in adult mice fed technical PCP-contaminated diets. Toxicol. Appl. Pharmacol. **62**:55–64.

KILZER, L., SCHEUNERT, I., GEYER, H., KLEIN, W. and KORTE, F. (1979). Laboratory screening of the volatilization rate of organic chemicals from water and soil. Chemosphere **8**:751–761.

KIMBROUGH, R.D. and LINDER, R.E. (1978). The effect of technical and purified pentachlorophenol on the rat liver. Toxicol. Appl. Pharmacol. **46**:151–162.

KIRSCH, E.J. and ETZEL, J.E. (1973). Microbial decomposition of pentachlorophenol. J. Water Pollut. Control Fed. **45**:359–364.

KLECKA, G.M. and MAIER, W.J. (1985). Kinetics of microbial growth on pentachlorophenol. Appl. Environ. Microbiol. **49**:46–53.

KNUDSEN, I., VERSCHUUREN, H.G., DEN TONKELAAR, E.M., KROES, R. and HELLEMAN, P.F.W. (1974). Short-term toxicity of pentachlorophenol in rats. Toxicology **2**:141–152.

KOBAYASHI, K. and AKITAKE, H. (1975). Studies on the metabolism of chlorophenols in fish. I. Absorption and excretion of PCP by goldfish. Bull. Jap. Soc. Sci. Fish. **41**:87–92. (Cited in U.S. EPA, 1980a).

KOBAYASHI, K. and KISHINO, T. (1980). Effect of pH on the toxicity and accumulation of pentachlorophenol in goldfish (*Carassius auratus*). Bull. Jap. Soc. Sci. Fish. **46**:167–170.

KOBAYASHI, K. and NAKAMURA, N. (1979). Studies on the metabolism of chlorophenols in fish. XII. Major detoxification pathways for pentachlorophenol in goldfish. Nippon Suisan Gakkaishi **45**:1085–1088. CA **92**:16544r.

KORTE, F. and KLEIN, W. (1982). Degradation of benzene in the environment. Ecotoxicol. Environ. Saf. **6**:311–327.

KORTE, F. and PARLAR, H. (1977). Decomposition of chlorinated hydrocarbons under simulated atmospheric conditions. In: Int. Conf. on Photochemical Oxidant Pollution and Its Control, pp. 729–736. U.S. EPA, Research Triangle Park, NC. EPA 600/3-77-001B.

KOTZIAS, D., PARLAR, H. and KORTE, F. (1982). Photoreactivity of organic chemicals in water in the presence of nitrate and nitrite. Naturwissen-schaften **69**:444–445.

KUEHL, D.W., LEONARD, E.N., WELCH, K.J. and VEITH, G.D. (1980). Identification of hazardous organic chemicals in fish from the Ashtabula River, Ohio and Wabash River, Indiana (USA). J. Assoc. Off. Anal. Chem. **63**:1238–1244.

KUEHL, D.W., LEONARD, E.N., BUTTERWORTH, B.C. and JOHNSON, K.L. (1983). Polychlorinated chemical residues in fish from major watersheds near the Great Lakes, 1979. Environ. Int. **9**:293–299.

KWASNIEWSKA, K. (1981). The degradation of pentachlorophenol by a fungus of *Fusarium* sp. Trace Subst. Environ. Health **15**:392–398. CA **97**(16):133081b.

LAMPARSKI, L.L., STEHL, R.H. and JOHNSON, R.L. (1980). Photolysis of pentachlorophenol-treated wood. Chlorinated dibenzo-p-dioxin formation. Environ. Sci. Technol. **14**:196–200.

LARSEN, R.V. (1976). The placental transfer and teratology of pentachlorophenol in rats. Diss. Abstr. Int. B **37**:1184–1185.

LARSEN, R.V., KIRSCH, L.E., SHAW, S.M., CHRISTIAN, J.E., JOHN, E. and BORN, G.S. (1972). Excretion and tissue distribution of uniformly labeled [^{14}C]-pentachlorophenol in rats. J. Pharmacol. Sci. **61**:2004–2006.

LARSEN, R.V., BORN, G.S., KESSLER, W.V., SHAW, S.M. and VAN SICKLE, D.C. (1975). Placental transfer and teratology of pentachlorophenol in rats. Environ. Lett. **10**:121–128. (Cited in U.S. EPA, 1982).

LEMMA, A. and AMES, B.N. (1975). Screening for mutagenic activity of some molluscicides. Trans. R. Soc. Trop. Med. Hyg. **69**:167–168. (Cited in U.S. EPA, 1984a).

LU, P.Y. and METCALF, R.L. (1975). Environmental fate and biodegradability of benzene derivates as studies in a model aquatic ecosystem. Environ. Health Perspect. **10**:269–284.

LYMAN, W.J., REEHE, W.F. and ROSENBLATT, D.H. (1982). Handbook of Chemical Property Estimation Methods, Environmental Behavior of Organic Compounds. McGraw-Hill, New York.

MABEY, W.R., SMITH, J.H., PODOLL, R.T. et al. (1981). Aquatic Fate Process Data for Organic Priority Pollutants. Monitoring and Data Support Div., U.S. EPA, Washington, DC. EPA 440/4-81-014.

MADDEN, M. and TITTLEBAUM, M. (1983). Biological treatability of pentachlorophenol wastewater. J. Environ. Sci. Health **A18**:583–596.

MARTIN, R.D. and WORTHING, C.R. (1977). Pesticide Manual, 5th ed., p. 403. British Crop Protection Council.

MELLO, W. (1978). National organics monitoring survey, March 1976 through January 1977. In: Investigations of Selected Environmental Pollutants. 1,2-Dichloroethane. EPA/560/2-78-006.

MIDDLETOWN, A.C. (1984). Land disposal and spill site environments. Basic Life Sci. **28**:137–149.

MITRUKA, B.M., ROWNSLEY, H.M. and VADEHA, D.V. (1976). Animals for medical research. In: Models for the Study of Human Disease. John Wiley, New York.

MOOS, L.P., KIRSCH, E.J., WUKASCH, R.F. and GRADY, C.P.L., JR. (1983). Pentachlorophenol. biodegradation. I. Aerobic Water Res. **17**:1575–1584.

MORIYA, M., OHTA, T., WATANABE, K., MIYAZAWA, T., KATO, K. and SHIRASU, Y. (1983). Further mutagenicity studies on pesticides in bacterial reversion assay system. Mutat. Res. **116**:185–216.

MURRAY, H.E., NEFF, G.S., HRUNG, Y. and GIAM, C.S. (1980). Determination of benzo(a)pyrene, hexachlorobenzene, and pentachlorophenol in oysters from Galveston Bay, TX. Bull. Environ. Contam. Toxicol. **25**:663–667.

MURRAY, H.E., RAY, L.E. and GIAM, C.S. (1981). Analysis of murine sediment, water and biota for selected organic pollutants. Chemosphere **10**:1327–1334.

MURTHY, N.B.K. and KAUFMAN, D.D. (1978). Degradation of pentachloronitrobenzene (PCNB) in anaerobic soils. J. Agric. Food Chem. **26**:1151–1156.

MURTHY, N.B.K., KAUFMAN, D.D. and FRIES, G.F. (1979). Degradation of pentachlorophenol (PCP) in aerobic and anaerobic soil. J. Environ. Sci. Health B. **15**:1–14.

NEW BRUNSWICK RES. PROD. COUNCIL. (1978). Bioaccumulation of toxic compounds in pulpmill effluents by aquatic organisms in receiving waters. CPAR Project Report 675-1. New Brunswick Research and Productivity Council, Fredericton, New Brunswick.

NIIMI, A.J. and CHO, C.Y. (1983). Laboratory and field analysis of pentachlorophenol (PCP) accumulation by salmonids. Water Res. **17**:1791–1795.

NIIMI, A.J. and MC FADDEN, C.A. (1982). Uptake of sodium pentachlorophenate (NaPCP) from water by rainbow trout (*Salmo gairdneri*) exposed to concentrations in the ng/L range. Bull. Environ. Contam. Toxicol. **28**:11–19.

NRC (National Research Council). (1977). Drinking Water and Health. National Academy of Science, Washington, DC. (Cited in U.S. EPA, 1980a).

NTP (National Toxicology Program) (1985). Management Status Report. November 5, 1985.

OBERG, T., AITTOLA, J.P. and BERGSTROEM, J.G.T. (1985). Chlorinated aromatics from the combustion of hazardous waste. Chemosphere **14**:215–221.

PAASIVIRTA, J., SARKKA, J., LESKIJARVI, T. and ROOS, A. (1980). Transportation and enrichment of chlorinated phenolic compounds in different aquatic food chains. Chemosphere **9**:441–456.

PAASIVIRTA, J., HEINOLA, K., HUMPPI, T. et al. (1985). Polychlorinated phenols, guaiacols and catecols in the environment. Chemosphere **14**:469–491.

PARRISH, P.R., DYAR, E.E., ENOS, J.M. and WILSON, W.G. (1978). Chronic toxicity of chlordane, trifluralin, and pentachlorophenol to sheepshead minnows (*Cyprinodon variegatus*). U.S. EPA, ORD. EPA 600/3-78-010.

PHIPPS, G.L. and HOLCOMBE, G.W. (1985). A method for aquatic multiple species toxicant testing: Acute toxicity of 10 chemicals to 5 vertebrates and 2 invertebrates. Environ. Pollut. Ser. A **38**:141–157.

PIERCE, R.H., JR. (1978). Fate and impact of pentachlorophenol in a freshwater ecosystem. EPA 600/3-78/063.

PIERCE, R.H., JR. and VICTOR, D.M. (1978). The fate of pentachlorophenol in an aquatic ecosystem. In: Chemistry, Pharmacology and Environmental Toxicology, K. R. Rao, ed., pp. 41–52. Plenum Press, New York.

PIERCE, R.H., JR., GOWER, S.A. and VICTOR, D.M. (1980). Pentachlorophenol and degradation products in lake sediment. Contam. Sediments **2**:43–56.

PIGNATELLO, J.J., MARTINSON, M.M., STEIERT, R.E., CARLSON, R.E. and CRAWFORD, R.L. (1983). Biodegradation and photolysis of pentachlorophenol in artificial freshwater streams. Appl. Environ. Microbiol. **46**:1024–1031.

PRUITT, G.W., GRANTHAJ, B.J. and PIERCE, R.H., JR. (1977). Accumulation and elimination of pentachlorophenol by the bluegill, *Lepomis macrochirus*. Trans. Am. Fish. Soc. **106**:462–465.

RAY, L.E., MURRAY, H.E. and GIAM, C.S. (1983a). Analysis of water and sediment from the Nueces Estuary/Corpus Christi Bay (Texas) for selected organic pollutants. Chemosphere **17**:1039–1045.

RAY, L.E., MURRAY, H.E. and GIAM, C.S. (1983b). Organic pollutants in marine samples from Portland, Maine. Chemosphere **12**:1031–1038.

RENBERG, L., MARTELL, E., SUNDSTROM, G. and ADOLFSSON-ERICI, M. (1983). Levels of chlorophenols in natural waters and fish after an accidental discharge of a wood-impregnating solution. Ambio. **12**:121–123.

SADTLER. (1966). Sadtler Standard Spectra. Sadtler Research Lab., Philadelphia, PA.

SANBORN, J.R., FRANCIS, B.M. and METCALF, R.L. (1977). The degradation of selected pesticides in soil: A review of published literature. U.S. EPA, Cincinnati, OH. EPA 600/9-77-022.

SCHAUERTE, W., LAY, J.P., KLEIN, W. and KORTE, F. (1982). Influence of 2,4,6-trichlorophenol and pentachlorophenol on the biota of aquatic systems. Outdoor experiments in compartments of a natural pond. Chemosphere **11**:71–79.

SCHELLENBERG, K., LEUNBERGER, C. and SCHWARZENBACH, R.P. (1984). Sorption of chlorinated phenols by natural sediments and aquifer materials. Environ. Sci. Technol. **18**:652–657.

SCHIMMEL, S.C., PATRICK, J.M., JR. and FAAS, L.F. (1978). Effects of sodium pentachlorophenate on several estuarine animals: Toxicity, uptake, and depuration. In: Chemistry, Pharmacology and Environmental Toxicology, K. R. Rao, ed., pp. 147–155. Plenum Press, New York.

SCHWETZ, B.A., KEELER, P.A. and GEHRING, P.J. (1974). Effect of purified and commercial grade pentachlorophenol on rat embryonal and fetal development. Toxicol. Appl. Pharmacol. **28**:151–161.

SCHWETZ, B.A., QUAST, J.F., KEELER, P.A., HUMISTON, C.G. and KOCIBA, R.J. (1977). Results of two-year toxicity and reproduction studies on pentachlorophenol in rats. In: Pentachlorophenol: Chemistry, Pharmacology and Environmental Toxicology, K. R. Rao, ed., pp. 301–309. Plenum Press, New York.

SKOW, K., GOYER, M., PAYNE, E. et al. (1980). An exposure and risk assessment for pentachlorophenol. U.S. EPA, Office of Water Regulations and Standards, Washington, DC. EPA 440/4-81-021.

SHACKELFORD, W.M. and KEITH, L.H. (1976). Frequency of organic compounds identified in water, pp. 190–191. U.S. EPA, Athens, GA. EPA 600/4-76-062.

SHIRASU, Y. (1976). J. Environ. Pollut. Control Fed. **12**:408. (Cited in U.S. EPA, 1984a).

SIMMON, V.F., KAUHANEN, K. and TARDIFF, R.G. (1977). Mutagenic activity of chemicals identified in drinking water. In: 2nd Intl. Conf. Environ. Mut.

SLOOF, F.W. (1983). Benthic macroinvertebrates and water quality assessment: Some toxicological considerations. Aquat. Toxicol. **4**:73–82.

SPEHAR, R.L., LEMKE, A.E., PICKERING, Q.H., ROUSCH, J.H., RUSSO, R.C. and YOUNT, J.D. (1981). Effects of pollution on freshwater fish. J. Water Pollut. Control Fed. **53**:1028–1076.

SPEHAR, R.L., NELSON, H.P., SWANSON, M.J. and RENOOS, J.W. (1985). Pentachlorophenol toxicity to amphipods and fathead minnows at different test pH values. Environ. Toxicol. Chem. **4**:389–397.

SRI (Stanford Research Institute). (1985). 1985 Directory of Chemical Producers: United States of America, p. 767. SRI International, Menlo Park, CA.

STATHAM, C.N., MELANCON, M.J., JR. and LECH, J.J. (1976). Bioconcentration of xenobiotics in trout bile: A proposed monitoring aid for some waterborne chemicals. Science **193**:680–681.

SUGIURA, K., AOKI, M., KENEKO, K. et al. (1984). Fate of 2,4,6-trichlorophenol, pentachlorophenol, p-chlorobiphenyl and hexachlorobenzene in an outdoor experimental pond: Comparison between observations and predictions based on laboratory data. Arch. Environ. Contam. Toxicol. **13**:745–758.

SUZUKI, T. and NOSE, K. (1970). Decomposition of pentachlorophenol in farm soil. Noyaki Seisan Gijutsu **22**:27–30. CA **76**:13140q.

TABAK, H.H., GUAVE, S.A., MASHNI, C.I. and BARTH, E.F. (1981). Biodegradability studies with organic priority pollutant compounds. J. Water Pollut. Control Fed. **53**:1503–1518.

TAGATZ, M.E., IVEY, J.M. and TOBIA, M. (1977). Effects of pentachlorophenol on development of estuarine communities. J. Toxicol. Environ. Health **3**:501–506.

TAGATZ, M.E., IVEY, J.M. and TOBIA, M. (1978). Effects of dowicide G-ST on development of experimental estuarine macrobenthic communities. In: Pentachlorophenol: Chemistry, Pharmacology and Environmental Toxicology, K. R. Rao, ed., pp. 157–163. Plenum Press, New York.

TAGATZ, M.E., IVEY, J.M., GREGORY, N.R. and OBLESBY, J.L. (1981). Effects of pentachlorophenol on field- and laboratory-developed estuarine benthic communities. Bull. Environ. Contam. Toxicol. **26**:137–143.

TREVORS, J.T. (1982). Effect of temperature on the biodegradation of pentachlorophenol by *Pseudomonas* species. Chemosphere **11**:471–475.

TRUJILLO, D.A., RAY, L.R., MURRAY, H.E. and GIAM, C.S. (1982). Bioaccumulation of pentachlorophenol by killfish (*Fundulus similus*). Chemosphere **11**:25–31.

U.S. EPA. (1978). Notice of Rebuttable Presumption against Registration and Continued Registration of Pesticide Products Containing Pentachlorophenol. Position Document 1. Federal Register **43**(202):48443–48477. Also available as EPA SPRD-80/85 and NTIS PB81-109464.

U.S. EPA. (1980a). Ambient Water Quality Criteria for Pentachlorophenol. Environmental Criteria and Assessment Office, Cincinnati, OH NTIS PB 81-117764.

U.S. EPA. (1980b). Guidelines and Methodology for the Preparation of the Health Effects Assessment Chapters of the Ambient Water Quality Criteria Documents. ECAO, OHEA, ORD, Cincinnati, OH. Also published in Fed. Reg. **45**:79318, 1980.

U.S. EPA. (1981a). Treatability Manual I. Treatability Data. U.S. EPA, Washington, DC. EPA-600/2-82-001A.

U.S. EPA. (1981b). Creosote, pentachlorophenol and inorganic arsenics: Preliminary Notice of Determination Concluding the RPAR of the Wood Preservative Uses of Pesticide Products; Notice of Availability of Position Document 2/3. Federal Register **46**(33):13020–13036.

U.S. EPA. (1982). Wood Preservative Pesticides Creosote, Pentachlorophenol and the Inorganic Arsenicals (Wood Uses) Position Document 2/3. U.S. EPA, Washington, DC, Office of Pesticides and Toxic Substances. EPA 540/9-82-004.

U.S. EPA (1984a). Drinking Water Criteria Document for Pentachlorophenol. Preliminary Program Office Draft. Prepared for Office of Drinking Water by ECAO, Cincinnati, OH. NTIS PB86-118015/AS.

U.S. EPA. (1984b). Proposed Guidelines for Carcinogen Risk Assessment. Federal Register **49**(227):46294–46301.

U.S. EPA. (1985a). STORET Water Quality Data Base. Online: October 1985.

U.S. EPA. (1985b). Summary of Acceptable Daily Intakes for Oral Exposure. Prepared by ECAO, Cincinnati, OH.

U.S. EPA. (1985c). National Primary Drinking Water Regulations: Synthetic organic chemicals, inorganic chemicals and microorganisms; proposed rule. 40 CFR **141**:46936–47025.

VALO, R., KITUNEN, V., SALKINOJA-SALONEN, M. and RAISANEN, S. (1984). Chlorinated phenols as contaminants of soil and water in the vicinity of two Finnish sawmills. Chemosphere **13**:835–844.

VAN DIJK, J.J., VAN DER MEER, C. and WIJNANS, M. (1977). The toxicity of sodium pentachlorophenate for three species of decapod crustaceans and their larvae. Bull. Environ. Contam. Toxicol. **17**:622–630.

VEITH, G.D., DE LORE, D.L. and BERGSTEDT, B.V. (1979a). Measuring and estimating the bioconcentration factor of chemicals in fish. J. Fish. Res. Board Can. **36**:1040–1048.

VEITH, G.D., KUEHL, D.W., LEONARD, E.N., PUGLISIS, F.A. and LEMKE, A.E. (1979b). Polychlorinated biphenyls and other organic chemical residues in fish from major watersheds of the United States, 1976. Pestic. Monit. J. **13**:1–11.

VERMA, S.R., RANI, S., TYAGI, A.K. and DALELA, R.C. (1980). Evaluation of acute toxicity of phenol and its chloro- and nitro-derivatives to certain teleosts. Water Air Soil Pollut. **14**:95–102.

VERMA, S.R., TONK, I.P. and DALELA, R.C. (1981). Determination of the maximum acceptable toxicant concentration (MATC) and the safe concentration for certain aquatic pollutants. Acta Hydrochim. Hydrobiol. **9**:247–254.

VERSCHUEREN, K. (1983). Handbook of Environmental Data on Organic Chemicals, 2nd ed., pp. 953–955. Van Nostrand Reinhold, New York.

VOGEL, E. and CHANDLER, J.L.R. (1974). Mutagenicity testing of cyclamate and some pesticides in *Drosophila melanogaster*. Experientia **30**:621–623.

WATANABE, I. (1973). Isolation of pentachlorophenol-degrading bacteria from soil. Soil Sci. Plant Nutr. **19**:109–116.

WATANABE, I., KASHIMOTO, T. and TATSUKAWA, R. (1985). Brominated phenols and anisoles in river and marine sediments in Japan. Bull. Environ. Contam. Toxicol. **35**:272–278.

WATERS, M.D., SANDHU, S.S., SIMMON, V.F. et al. (1982). Study of pesticide genotoxicity. Basic Life Sci. **21**:275–326.

WEBB, P.W. and BRETT, J.R. (1973). Effects of sublethal concentrations of sodium pentachlorophenate on growth rate, food conversion efficiency, and swimming performance in underyearling sockeye salmon (*Oncorhynchus nerka*). J. Fish. Res. Board Can. **30**:499.

WEGMAN, R.C.C. and VANDENBROEK, H.H. (1983). Chlorophenols in river sediment in The Netherlands. Water Res. **17**:227–230.

WEISS, U.M., SCHEUNERT, I., KLEIN, W. and KORTE, F. (1982). Fate of pentachlorophenol-C14 in soil under controlled conditions. J. Agric. Food Chem. **30**:1191–1194.

WONG, A.S. and CROSBY, D.G. (1981). Photodecomposition of pentachlorophenol in water. J. Agric. Food Chem. **29**:125–130.

WYLLIE, J.A., GABICA, J., BENSON, W.W. and YODER, J. (1975). Exposure and contamination of the air and employees of a pentachlorophenol plant, Idaho-1972. Pestic. Monit. J. **9**:150–153.

XIE, T.M. (1983). Determination of trace amounts on chlorophenols and chloroguaiacols in sediment. Chemosphere **12**:1183–1192.

YOUNG, D.R., GOSSET, R.W., BAIRD, R.B., BROWN, D.A., TAYLOR, P.A. and MIILLE, M.J. (1983). Wastewater inputs and marine bioaccumulation of priority pollutant organics off southern California. Water Chlorination: Environ. Impact Health Eff. **4**:871–884.

ZEPP, R.G., SCHLOTZHAUER, P.F., SIMMONS, M.S., MILLER, G.C., BAUGHMAN, G.L. and WOLFE, N.L. (1984). Dynamics of pollutant photoreactions in the hydrosphere. Fresenius Z. Anal. Chem. **319**:119–125.

ZITKO, V., HUTZINGER, O. and CHOI, P.M.K. (1974). Determination of pentachlorophenol and chlorobiphenyls in biological samples. Bull. Environ. Contam. Toxicol. **12**:649–653.

APPENDIX
Literature Searched

This profile is based on data identified by computerized literature searches of the following:

CASR online (U.S. EPA Chemical Activities Status Report)
CAS online STN International
TOXLINE
TOXBACK 76
TOXBACK 65
RTECS
OHM TADS
STORET
SRC Environmental Fate Data Bases
SANSS
AQUIRE
TSCAPP
NTIS
Federal Register.

These searches were conducted in October 1985. In addition, hand searches were made of Chemical Abstracts (Collective Indices 6 and 7), and the following secondary sources were reviewed:

ACGIH (American Conference of Governmental Industrial Hygienists). (1980). Documentation of the Threshold Limit Values, 4th ed. (Includes Supplemental Documentation, 1981, 1982, 1983). Cincinnati, OH.

ACHIH. (1985). TLVs: Threshold Limit Values for Chemical Substances and Physical Agents in the Workroom Environment with Intended Changes for 1985-1986. Cincinnati, OH.

CLAYTON, G.D. and CLAYTON, F.E., eds. (1981). Patty's Industrial Hygiene and Toxicology, 3rd rev. ed., Vol. 2A. John Wiley, New York.

CLAYTON, G.D. and CLAYTON, F.E., eds. (1981). Patty's Industrial Industrial Hygiene and Toxicology, 3rd rev. ed., Vol. 2B. John Wiley, New York.

CLAYTON, G.D. and CLAYTON F.E., eds. (1982). Patty's Industrial Hygiene and Toxicology, 3rd rev. ed., Vol. 2C. John Wiley, New York.

GRAYSON, M. and ECKROTH, D., eds. (1978-1983). Kirk-Othmer Encyclopedia of Chemical Technology, 3rd ed., 23 vols. John Wiley, New York.

HAMILTON, A. and HARDY, H.L. (1974). Industrial Toxicology, 3rd ed. Publishing Sciences Group, Boston, MA.

IARC (International Agency for Research on Cancer). IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans. WHO, IARC, Lyons, France.

ITII (International Technical Information Institute). (1982). Toxic and Hazardous Industrial Chemicals Safety Manual for Handling and Disposal with Toxicity and Hazard Data. ITII, Tokyo, Japan.

NTP (National Toxicology Program). (1984). Toxicology Research and Testing Program. Chemicals on Standard Protocol. Management Status.

OUELLETTE, R.P. and KING, J.A. (1977). Chemical Week Pesticide Register. McGraw-Hill, New York.

SAX, N.I. (1979). Dangerous Properties of Industrial Materials, 5th ed. Van Nostrand Reinhold, New York.

SRI (Stanford Research Institute). (1984). Directory of Chemical Producers. Menlo Park, CA.

U.S. EPA (United States Environmental Protection Agency). (1985). Status Report on Rebuttable Presumption against Registration (RPAR) or Special Review Process. Registration Standards and the Data Call-in Programs. Office of Pesticide Programs, Washington, D.C.

U.S. EPA. (1985). CSB Existing Chemical Assessment Tracking System. Name and CAS Number Ordered Indexes. Office of Toxic Substances, Washington, DC.

USITC (U.S. International Trade Commission). (1983). Synthetic Organic Chemicals. U.S. Production and Sales, 1982. USITC Publ. 1422, Washington, DC.

VERSCHUEREN, K. (1983). Handbook of Environmental Data on Organic Chemicals, 2nd ed. Van Nostrand Reinhold, New York.

WORTHING, C.R. and WALKER, S.B., eds. (1983). The Pesticide Manual. British Crop Protection Council.

WINDHOLTZ, M., ed. (1983). The Merck Index, 10th ed. Merck and Co., Inc., Rahway, NJ.

In addition, approximately 30 compendia of aquatic toxicity data were reviewed, including the following:

BATTELLE'S COLUMBUS LABORATORIES. (1971). Water Quality Criteria Data Book. Volume 3. Effects of Chemicals on Aquatic Life. Selected Data from the Literature through 1968. Prepared for the U.S. EPA under Contract No. 68-01-0007. Washington, DC.

JOHNSON, W.W. and FINLEY, M.T. (1980). Handbook of Acute Toxicity of Chemicals to Fish and Aquatic Invertebrates. Summaries of the Toxicity Tests Conducted at Columbia National Fisheries Research Laboratory, 1965-1978. U.S. Dept. Interior, Fish and Wildlife Serv. Res. Publ. 137, Washington, DC.

MC FEE, J.D. and WOLF, H.W. (1963). Water Quality Criteria, 2nd ed. Prepared for the Resources Agency of California, State Water Quality Control Board. Publ. No. 3-A.

PIMENTAL, D. (1971). Ecological Effects of Pesticides on Non-Target Species. Prepared for the U.S. EPA, Washington, DC. PB-269605.

SCHNEIDER, B.A. (1979). Toxicology Handbook. Mammalian and Aquatic Data. Book 1: Toxicology Data. Office of Pesticide Programs, U.S. EPA, Washington, DC. EPA 540/9-79-003. NTIS PB 80-196876,

Received September 12, 1986
Accepted October 17, 1986

NOTICE

This document has been reviewed in accordance with U.S. Environmental Protection Agency policy and approved for publication. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

This article documents research that was conducted under the auspices of the U.S. Environmental Protection Agency (USEPA). As an arm of the government, the USEPA has a paid up, non-exclusive, irrevocable, worldwide license to reproduce, prepare derivative works, and distribute to the public all of the material contained herein, and to grant others acting on its behalf the right to do so. The USEPA grants to the National Technical Information Service, Department of Commerce, the right to act as its agent in the distribution of this material.